

# Collins asymmetries in inclusive charged $KK$ and $K\pi$ pairs produced in $e^+e^-$ annihilation

J. P. Lees,<sup>1</sup> V. Poireau,<sup>1</sup> V. Tisserand,<sup>1</sup> E. Grauges,<sup>2</sup> A. Palano,<sup>3a,3b</sup> G. Eigen,<sup>4</sup> B. Stugu,<sup>4</sup> D. N. Brown,<sup>5</sup> L. T. Kerth,<sup>5</sup> Yu. G. Kolomensky,<sup>5</sup> M. J. Lee,<sup>5</sup> G. Lynch,<sup>5</sup> H. Koch,<sup>6</sup> T. Schroeder,<sup>6</sup> C. Hearty,<sup>7</sup> T. S. Mattison,<sup>7</sup> J. A. McKenna,<sup>7</sup> R. Y. So,<sup>7</sup> A. Khan,<sup>8</sup> V. E. Blinov,<sup>9a,9b,9c</sup> A. R. Buzykaev,<sup>9a</sup> V. P. Druzhinin,<sup>9a,9b</sup> V. B. Golubev,<sup>9a,9b</sup> E. A. Kravchenko,<sup>9a,9b</sup> A. P. Onuchin,<sup>9a,9b,9c</sup> S. I. Serednyakov,<sup>9a,9b</sup> Yu. I. Skovpen,<sup>9a,9b</sup> E. P. Solodov,<sup>9a,9b</sup> K. Yu. Todyshev,<sup>9a,9b</sup> A. J. Lankford,<sup>10</sup> B. Dey,<sup>11</sup> J. W. Gary,<sup>11</sup> O. Long,<sup>11</sup> M. Franco Sevilla,<sup>12</sup> T. M. Hong,<sup>12</sup> D. Kovalskyi,<sup>12</sup> J. D. Richman,<sup>12</sup> C. A. West,<sup>12</sup> A. M. Eisner,<sup>13</sup> W. S. Lockman,<sup>13</sup> W. Panduro Vazquez,<sup>13</sup> B. A. Schumm,<sup>13</sup> A. Seiden,<sup>13</sup> D. S. Chao,<sup>14</sup> C. H. Cheng,<sup>14</sup> B. Echenard,<sup>14</sup> K. T. Flood,<sup>14</sup> D. G. Hitlin,<sup>14</sup> T. S. Miyashita,<sup>14</sup> P. Ongmongkolkul,<sup>14</sup> F. C. Porter,<sup>14</sup> M. Röhrken,<sup>14</sup> R. Andreassen,<sup>15</sup> Z. Huard,<sup>15</sup> B. T. Meadows,<sup>15</sup> B. G. Pushpawela,<sup>15</sup> M. D. Sokoloff,<sup>15</sup> L. Sun,<sup>15</sup> P. C. Bloom,<sup>16</sup> W. T. Ford,<sup>16</sup> A. Gaz,<sup>16</sup> J. G. Smith,<sup>16</sup> S. R. Wagner,<sup>16</sup> R. Ayad,<sup>17,†</sup> W. H. Toki,<sup>17</sup> B. Spaan,<sup>18</sup> D. Bernard,<sup>19</sup> M. Verderi,<sup>19</sup> S. Playfer,<sup>20</sup> D. Bettoni,<sup>21a</sup> C. Bozzi,<sup>21a</sup> R. Calabrese,<sup>21a,21b</sup> G. Cibinetto,<sup>21a,21b</sup> E. Fioravanti,<sup>21a,21b</sup> I. Garzia,<sup>21a,21b</sup> E. Luppi,<sup>21a,21b</sup> L. Piemontese,<sup>21a</sup> V. Santoro,<sup>21a</sup> A. Calcaterra,<sup>22</sup> R. de Sangro,<sup>22</sup> G. Finocchiaro,<sup>22</sup> S. Martellotti,<sup>22</sup> P. Patteri,<sup>22</sup> I. M. Peruzzi,<sup>22,‡</sup> M. Piccolo,<sup>22</sup> A. Zallo,<sup>22</sup> R. Contri,<sup>23a,23b</sup> M. R. Monge,<sup>23a,23b</sup> S. Passaggio,<sup>23a</sup> C. Patrignani,<sup>23a</sup> B. Bhuyan,<sup>24</sup> V. Prasad,<sup>24</sup> A. Adametz,<sup>25</sup> U. Uwer,<sup>25</sup> H. M. Lacker,<sup>26</sup> U. Mallik,<sup>27</sup> C. Chen,<sup>28</sup> J. Cochran,<sup>28</sup> S. Prell,<sup>28</sup> H. Ahmed,<sup>29</sup> A. V. Gritsan,<sup>30</sup> N. Arnaud,<sup>31</sup> M. Davier,<sup>31</sup> D. Derkach,<sup>31</sup> G. Grosdidier,<sup>31</sup> F. Le Diberder,<sup>31</sup> A. M. Lutz,<sup>31</sup> B. Malaescu,<sup>31,§</sup> P. Roudeau,<sup>31</sup> A. Stocchi,<sup>31</sup> G. Wormser,<sup>31</sup> D. J. Lange,<sup>32</sup> D. M. Wright,<sup>32</sup> J. P. Coleman,<sup>33</sup> J. R. Fry,<sup>33</sup> E. Gabathuler,<sup>33</sup> D. E. Hutchcroft,<sup>33</sup> D. J. Payne,<sup>33</sup> C. Touramanis,<sup>33</sup> A. J. Bevan,<sup>34</sup> F. Di Lodovico,<sup>34</sup> R. Sacco,<sup>34</sup> G. Cowan,<sup>35</sup> D. N. Brown,<sup>36</sup> C. L. Davis,<sup>36</sup> A. G. Denig,<sup>37</sup> M. Fritsch,<sup>37</sup> W. Gradl,<sup>37</sup> K. Griessinger,<sup>37</sup> A. Hafner,<sup>37</sup> K. R. Schubert,<sup>37</sup> R. J. Barlow,<sup>38,¶</sup> G. D. Lafferty,<sup>38</sup> R. Cenci,<sup>39</sup> B. Hamilton,<sup>39</sup> A. Jawahery,<sup>39</sup> D. A. Roberts,<sup>39</sup> R. Cowan,<sup>40</sup> R. Cheaib,<sup>41</sup> P. M. Patel,<sup>41,\*</sup> S. H. Robertson,<sup>41</sup> N. Neri,<sup>42a</sup> F. Palombo,<sup>42a,42b</sup> L. Cremaldi,<sup>43</sup> R. Godang,<sup>43,\*\*</sup> D. J. Summers,<sup>43</sup> M. Simard,<sup>44</sup> P. Taras,<sup>44</sup> G. De Nardo,<sup>45a,45b</sup> G. Onorato,<sup>45a,45b</sup> C. Sciacca,<sup>45a,45b</sup> G. Raven,<sup>46</sup> C. P. Jessop,<sup>47</sup> J. M. LoSecco,<sup>47</sup> K. Honscheid,<sup>48</sup> R. Kass,<sup>48</sup> M. Margoni,<sup>49a,49b</sup> M. Morandin,<sup>49a</sup> M. Posocco,<sup>49a</sup> M. Rotondo,<sup>49a</sup> G. Simi,<sup>49a,49b</sup> F. Simonetto,<sup>49a,49b</sup> R. Stroili,<sup>49a,49b</sup> S. Akar,<sup>50</sup> E. Ben-Haim,<sup>50</sup> M. Bomben,<sup>50</sup> G. R. Bonneaud,<sup>50</sup> H. Briand,<sup>50</sup> G. Calderini,<sup>50</sup> J. Chauveau,<sup>50</sup> Ph. Leruste,<sup>50</sup> G. Marchiori,<sup>50</sup> J. Ocariz,<sup>50</sup> M. Biasini,<sup>51a,51b</sup> E. Manoni,<sup>51a</sup> A. Rossi,<sup>51a</sup> C. Angelini,<sup>52a,52b</sup> G. Batignani,<sup>52a,52b</sup> S. Bettarini,<sup>52a,52b</sup> M. Carpinelli,<sup>52a,52b,††</sup> G. Casarosa,<sup>52a,52b</sup> M. Chrzaszcz,<sup>52a</sup> F. Forti,<sup>52a,52b</sup> M. A. Giorgi,<sup>52a,52b</sup> A. Lusiani,<sup>52a,52c</sup> B. Oberhof,<sup>52a,52b</sup> E. Paoloni,<sup>52a,52b</sup> M. Rama,<sup>52a</sup> G. Rizzo,<sup>52a,52b</sup> J. J. Walsh,<sup>52a</sup> D. Lopes Pegna,<sup>53</sup> J. Olsen,<sup>53</sup> A. J. S. Smith,<sup>53</sup> F. Anulli,<sup>54a</sup> R. Faccini,<sup>54a,54b</sup> F. Ferrarotto,<sup>54a</sup> F. Ferroni,<sup>54a,54b</sup> M. Gaspero,<sup>54a,54b</sup> A. Pilloni,<sup>54a,54b</sup> G. Piredda,<sup>54a</sup> C. Büniger,<sup>55</sup> S. Dittrich,<sup>55</sup> O. Grünberg,<sup>55</sup> M. Hess,<sup>55</sup> T. Leddig,<sup>55</sup> C. Voß,<sup>55</sup> R. Waldi,<sup>55</sup> T. Adye,<sup>56</sup> E. O. Olaiya,<sup>56</sup> F. F. Wilson,<sup>56</sup> S. Emery,<sup>57</sup> G. Vasseur,<sup>57</sup> D. Aston,<sup>58</sup> D. J. Bard,<sup>58</sup> C. Cartaro,<sup>58</sup> M. R. Convery,<sup>58</sup> J. Dorfan,<sup>58</sup> G. P. Dubois-Felsmann,<sup>58</sup> W. Dunwoodie,<sup>58</sup> M. Ebert,<sup>58</sup> R. C. Field,<sup>58</sup> B. G. Fulsom,<sup>58</sup> M. T. Graham,<sup>58</sup> C. Hast,<sup>58</sup> W. R. Innes,<sup>58</sup> P. Kim,<sup>58</sup> D. W. G. S. Leith,<sup>58</sup> S. Luitz,<sup>58</sup> V. Luth,<sup>58</sup> D. B. MacFarlane,<sup>58</sup> D. R. Muller,<sup>58</sup> H. Neal,<sup>58</sup> T. Pulliam,<sup>58</sup> B. N. Ratcliff,<sup>58</sup> A. Roodman,<sup>58</sup> R. H. Schindler,<sup>58</sup> A. Snyder,<sup>58</sup> D. Su,<sup>58</sup> M. K. Sullivan,<sup>58</sup> J. Va'vra,<sup>58</sup> W. J. Wisniewski,<sup>58</sup> H. W. Wulsin,<sup>58</sup> M. V. Purohit,<sup>59</sup> J. R. Wilson,<sup>59</sup> A. Randle-Conde,<sup>60</sup> S. J. Sekula,<sup>60</sup> M. Bellis,<sup>61</sup> P. R. Burchat,<sup>61</sup> E. M. T. Puccio,<sup>61</sup> M. S. Alam,<sup>62</sup> J. A. Ernst,<sup>62</sup> R. Gorodeisky,<sup>63</sup> N. Guttman,<sup>63</sup> D. R. Peimer,<sup>63</sup> A. Soffer,<sup>63</sup> S. M. Spanier,<sup>64</sup> J. L. Ritchie,<sup>65</sup> R. F. Schwitters,<sup>65</sup> J. M. Izen,<sup>66</sup> X. C. Lou,<sup>66</sup> F. Bianchi,<sup>67a,67b</sup> F. De Mori,<sup>67a,67b</sup> A. Filippi,<sup>67a</sup> D. Gamba,<sup>67a,67b</sup> L. Lanceri,<sup>68a,68b</sup> L. Vitale,<sup>68a,68b</sup> F. Martinez-Vidal,<sup>69</sup> A. Oyanguren,<sup>69</sup> J. Albert,<sup>70</sup> Sw. Banerjee,<sup>70</sup> A. Beaulieu,<sup>70</sup> F. U. Bernlochner,<sup>70</sup> H. H. F. Choi,<sup>70</sup> G. J. King,<sup>70</sup> R. Kowalewski,<sup>70</sup> M. J. Lewczuk,<sup>70</sup> T. Lueck,<sup>70</sup> I. M. Nugent,<sup>70</sup> J. M. Roney,<sup>70</sup> R. J. Sobie,<sup>70</sup> N. Tasneem,<sup>70</sup> T. J. Gershon,<sup>71</sup> P. F. Harrison,<sup>71</sup> T. E. Latham,<sup>71</sup> H. R. Band,<sup>72</sup> S. Dasu,<sup>72</sup> Y. Pan,<sup>72</sup> R. Prepost,<sup>72</sup> and S. L. Wu<sup>72</sup>

(BABAR Collaboration)

<sup>1</sup>Laboratoire d'Annecy-le-Vieux de Physique des Particules (LAPP), Université de Savoie, CNRS/IN2P3, F-74941 Annecy-Le-Vieux, France

<sup>2</sup>Universitat de Barcelona, Facultat de Física, Departament ECM, E-08028 Barcelona, Spain

<sup>3a</sup>INFN Sezione di Bari, I-70126 Bari, Italy

<sup>3b</sup>Dipartimento di Fisica, Università di Bari, I-70126 Bari, Italy

<sup>4</sup>University of Bergen, Institute of Physics, N-5007 Bergen, Norway

<sup>5</sup>Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

<sup>6</sup>Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

<sup>7</sup>University of British Columbia, Vancouver, British Columbia V6T 1Z1, Canada

<sup>8</sup>Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

<sup>9a</sup>Budker Institute of Nuclear Physics, SB RAS, Novosibirsk 630090, Russia

<sup>9b</sup>Novosibirsk State University, Novosibirsk 630090, Russia

<sup>9c</sup>Novosibirsk State Technical University, Novosibirsk 630092, Russia

- <sup>10</sup>University of California at Irvine, Irvine, California 92697, USA
- <sup>11</sup>University of California at Riverside, Riverside, California 92521, USA
- <sup>12</sup>University of California at Santa Barbara, Santa Barbara, California 93106, USA
- <sup>13</sup>University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA
- <sup>14</sup>California Institute of Technology, Pasadena, California 91125, USA
- <sup>15</sup>University of Cincinnati, Cincinnati, Ohio 45221, USA
- <sup>16</sup>University of Colorado, Boulder, Colorado 80309, USA
- <sup>17</sup>Colorado State University, Fort Collins, Colorado 80523, USA
- <sup>18</sup>Technische Universität Dortmund, Fakultät Physik, D-44221 Dortmund, Germany
- <sup>19</sup>Laboratoire Leprince-Ringuet, Ecole Polytechnique, CNRS/IN2P3, F-91128 Palaiseau, France
- <sup>20</sup>University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom
- <sup>21a</sup>INFN Sezione di Ferrara, I-44122 Ferrara, Italy
- <sup>21b</sup>Dipartimento di Fisica e Scienze della Terra, Università di Ferrara, I-44122 Ferrara, Italy
- <sup>22</sup>INFN Laboratori Nazionali di Frascati, I-00044 Frascati, Italy
- <sup>23a</sup>INFN Sezione di Genova, I-16146 Genova, Italy
- <sup>23b</sup>Dipartimento di Fisica, Università di Genova, I-16146 Genova, Italy
- <sup>24</sup>Indian Institute of Technology Guwahati, Guwahati, Assam 781 039, India
- <sup>25</sup>Universität Heidelberg, Physikalisches Institut, D-69120 Heidelberg, Germany
- <sup>26</sup>Humboldt-Universität zu Berlin, Institut für Physik, D-12489 Berlin, Germany
- <sup>27</sup>University of Iowa, Iowa City, Iowa 52242, USA
- <sup>28</sup>Iowa State University, Ames, Iowa 50011-3160, USA
- <sup>29</sup>Physics Department, Jazan University, Jazan 22822, Saudi Arabia
- <sup>30</sup>Johns Hopkins University, Baltimore, Maryland 21218, USA
- <sup>31</sup>Laboratoire de l'Accélérateur Linéaire, IN2P3/CNRS et Université Paris-Sud 11, Centre Scientifique d'Orsay, F-91898 Orsay Cedex, France
- <sup>32</sup>Lawrence Livermore National Laboratory, Livermore, California 94550, USA
- <sup>33</sup>University of Liverpool, Liverpool L69 7ZE, United Kingdom
- <sup>34</sup>Queen Mary, University of London, London, E1 4NS, United Kingdom
- <sup>35</sup>University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom
- <sup>36</sup>University of Louisville, Louisville, Kentucky 40292, USA
- <sup>37</sup>Johannes Gutenberg-Universität Mainz, Institut für Kernphysik, D-55099 Mainz, Germany
- <sup>38</sup>University of Manchester, Manchester M13 9PL, United Kingdom
- <sup>39</sup>University of Maryland, College Park, Maryland 20742, USA
- <sup>40</sup>Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA
- <sup>41</sup>McGill University, Montréal, Québec, Canada H3A 2T8
- <sup>42a</sup>INFN Sezione di Milano, I-20133 Milano, Italy
- <sup>42b</sup>Dipartimento di Fisica, Università di Milano, I-20133 Milano, Italy
- <sup>43</sup>University of Mississippi, University, Mississippi 38677, USA
- <sup>44</sup>Université de Montréal, Physique des Particules, Montréal, Québec H3C 3J7, Canada
- <sup>45a</sup>INFN Sezione di Napoli, I-80126 Napoli, Italy
- <sup>45b</sup>Dipartimento di Scienze Fisiche, Università di Napoli Federico II, I-80126 Napoli, Italy
- <sup>46</sup>NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
- <sup>47</sup>University of Notre Dame, Notre Dame, Indiana 46556, USA
- <sup>48</sup>Ohio State University, Columbus, Ohio 43210, USA
- <sup>49a</sup>INFN Sezione di Padova, I-35131 Padova, Italy
- <sup>49b</sup>Dipartimento di Fisica, Università di Padova, I-35131 Padova, Italy
- <sup>50</sup>Laboratoire de Physique Nucléaire et de Hautes Energies, IN2P3/CNRS, Université Pierre et Marie Curie-Paris 6, Université Denis Diderot-Paris 7, F-75252 Paris, France
- <sup>51a</sup>INFN Sezione di Perugia, I-06123 Perugia, Italy
- <sup>51b</sup>Dipartimento di Fisica, Università di Perugia, I-06123 Perugia, Italy
- <sup>52a</sup>INFN Sezione di Pisa, I-56127 Pisa, Italy
- <sup>52b</sup>Dipartimento di Fisica, Università di Pisa, I-56127 Pisa, Italy
- <sup>52c</sup>Scuola Normale Superiore di Pisa, I-56127 Pisa, Italy
- <sup>53</sup>Princeton University, Princeton, New Jersey 08544, USA
- <sup>54a</sup>INFN Sezione di Roma, I-00185 Roma, Italy
- <sup>54b</sup>Dipartimento di Fisica, Università di Roma La Sapienza I-00185 Roma, Italy
- <sup>55</sup>Universität Rostock, D-18051 Rostock, Germany

- <sup>56</sup>Rutherford Appleton Laboratory, Chilton, Didcot, Oxon OX11 0QX, United Kingdom  
<sup>57</sup>CEA, Irfu, SPP, Centre de Saclay, F-91191 Gif-sur-Yvette, France  
<sup>58</sup>SLAC National Accelerator Laboratory, Stanford, California 94309 USA  
<sup>59</sup>University of South Carolina, Columbia, South Carolina 29208, USA  
<sup>60</sup>Southern Methodist University, Dallas, Texas 75275, USA  
<sup>61</sup>Stanford University, Stanford, California 94305-4060, USA  
<sup>62</sup>State University of New York, Albany, New York 12222, USA  
<sup>63</sup>Tel Aviv University, School of Physics and Astronomy, Tel Aviv 69978, Israel  
<sup>64</sup>University of Tennessee, Knoxville, Tennessee 37996, USA  
<sup>65</sup>University of Texas at Austin, Austin, Texas 78712, USA  
<sup>66</sup>University of Texas at Dallas, Richardson, Texas 75083, USA  
<sup>67a</sup>INFN Sezione di Torino, I-10125 Torino, Italy  
<sup>67b</sup>Dipartimento di Fisica, Università di Torino, I-10125 Torino, Italy  
<sup>68a</sup>INFN Sezione di Trieste, I-34127 Trieste, Italy  
<sup>68b</sup>Dipartimento di Fisica, Università di Trieste, I-34127 Trieste, Italy  
<sup>69</sup>IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain  
<sup>70</sup>University of Victoria, Victoria, British Columbia V8W 3P6, Canada  
<sup>71</sup>Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom  
<sup>72</sup>University of Wisconsin, Madison, Wisconsin 53706, USA  
(Received 23 June 2015; published 22 December 2015)

We present measurements of Collins asymmetries in the inclusive process  $e^+e^- \rightarrow h_1 h_2 X$ ,  $h_1 h_2 = KK$ ,  $K\pi$ ,  $\pi\pi$ , at the center-of-mass energy of 10.6 GeV, using a data sample of 468 fb<sup>-1</sup> collected by the BABAR experiment at the PEP-II  $B$  factory at SLAC National Accelerator Center. Considering hadrons in opposite thrust hemispheres of hadronic events, we observe clear azimuthal asymmetries in the ratio of unlike sign to like sign, and unlike sign to all charged  $h_1 h_2$  pairs, which increase with hadron energies. The  $K\pi$  asymmetries are similar to those measured for the  $\pi\pi$  pairs, whereas those measured for high-energy  $KK$  pairs are, in general, larger.

DOI: 10.1103/PhysRevD.92.111101

PACS numbers: 13.66.Bc, 13.87.Fh, 13.88.+e, 14.65.-q

The Collins effect [1] relates the transverse spin component of a fragmenting quark to the azimuthal distribution of final state hadrons about its flight direction. The chiral-odd, transverse momentum-dependent Collins fragmentation function (FF) provides a unique probe of QCD, such as factorization and evolution with the energy scale  $Q^2$  [2–5].

Additional interest has been sparked by the observation of azimuthal asymmetries for pions and kaons in semi-inclusive deep inelastic scattering experiments (SIDIS) [6–10]. These are sensitive to the product of a Collins FF and a chiral-odd transversity parton distribution function (PDF), one of the three fundamental PDFs needed to describe the spin content of the nucleon. Although these observations require nonzero Collins FFs, independent

direct measurements of one of these chiral-odd functions are needed to determine each of them.

In  $e^+e^-$  annihilation, one can measure the product of two Collins FFs, and detailed measurements have been made for pairs of charged pions [11–13]. No measurements are available for  $K\pi$  and  $KK$  pairs, which are sensitive to different quark-flavor combinations, in particular the contribution of the strange quark. Such measurements could be combined with SIDIS data to simultaneously determine the Collins FFs and transversity PDF for up, down, and strange quarks [14–19].

In this paper, we report the measurement of the Collins effect (or Collins asymmetry) for inclusive production of hadron pairs in the process  $e^+e^- \rightarrow q\bar{q} \rightarrow h_1 h_2 X$ , where  $h_{1,2} = K^\pm$  or  $\pi^\pm$ ,  $q$  stands for light quarks  $u$  or  $d$  or  $s$ , and  $X$  for any combination of additional hadrons.

The probability that a transversely polarized quark ( $q^\uparrow$ ) with momentum direction  $\hat{\mathbf{k}}$  and spin  $\mathbf{S}_q$  fragments into a hadron  $h$  carrying zero intrinsic spin with momentum  $\mathbf{P}_h$  is defined in terms of unpolarized  $D_1^q$  and Collins  $H_1^{\perp q}$  fragmentation functions [20]:

$$D_h^{q^\uparrow}(z, \mathbf{P}_{hT}) = D_1^q(z, P_{hT}^2) + H_1^{\perp q}(z, P_{hT}^2) \frac{(\hat{\mathbf{k}} \times \mathbf{P}_{hT}) \cdot \mathbf{S}_q}{zM_h}, \quad (1)$$

\*Deceased.

†Present address: University of Tabuk, Tabuk 71491, Saudi Arabia.

‡Also at Università di Perugia, Dipartimento di Fisica, I-06123 Perugia, Italy.

§Present address: Laboratoire de Physique Nucléaire et de Hautes Energies, IN2P3/CNRS, F-75252 Paris, France.

¶Present address: University of Huddersfield, Huddersfield HD1 3DH, UK.

\*\*Present address: University of South Alabama, Mobile, Alabama 36688, USA.

††Also at Università di Sassari, I-07100 Sassari, Italy.

where  $M_h$ ,  $\mathbf{P}_{hT}$ , and  $z = 2E_h/\sqrt{s}$  are the hadron mass, momentum transverse to  $\hat{\mathbf{k}}$ , and fractional energy, respectively, with  $E_h$  its total energy and  $\sqrt{s}$  the  $e^+e^-$  center-of-mass (c.m.) energy. The term including  $H_1^\perp$  introduces an azimuthal modulation around the direction of the fragmenting quark, called Collins asymmetry.

In  $e^+e^- \rightarrow q\bar{q}$  events, the  $q$  and  $\bar{q}$  must be produced back to back in the  $e^+e^-$  c.m. frame with their spin aligned. For unpolarized  $e^+$  and  $e^-$  beams at *BABAR* energies, the  $q$  and  $\bar{q}$  spins are polarized along either the  $e^+$  or  $e^-$  beam direction, so there is a large transverse component when the angle between the  $e^+e^-$  and the  $q\bar{q}$  axis is large. The direction is unknown for a given event, but the correlation can be exploited. Experimentally, the  $q$  and  $\bar{q}$  directions are difficult to measure, but the event thrust axis  $\hat{n}$  [21,22] approximates at leading order the  $q\bar{q}$  axis, so an azimuthal correlation between two hadrons in opposite thrust hemispheres reflects the product of the two Collins functions.

Figure 1 shows the thrust reference frame (RF12) [23]. If not otherwise specified, all kinematic variables are defined in the  $e^+e^-$  c.m. frame. The Collins effect results in a cosine modulation of the azimuthal angle  $\phi_{12} = \phi_1 + \phi_2$  of the dihadron yields. Expressing the yield as a function of  $\phi_{12}$  (after the integration over  $\mathbf{P}_{hT}$ ), and dividing by the average bin content, we obtain the normalized rate [11]

$$R_{12}(\phi_{12}) = 1 + \frac{\sin^2\theta_{th}}{1 + \cos^2\theta_{th}} \cos\phi_{12} \cdot \frac{H_1^{\perp[1]}(z_1)\bar{H}_1^{\perp[1]}(z_2)}{D_1^{[0]}(z_1)\bar{D}_1^{[0]}(z_2)}, \quad (2)$$

where the sum over the involved quark flavors is implied,  $\theta_{th}$  is defined in Fig. 1,  $z_{1(2)}$  is the fractional energy of the first (second) hadron, and the bar denotes the function for the  $\bar{q}$ . Equation (2) involves only the moments of FF, which are defined as




FIG. 1 (color online). Thrust reference frame (RF12). The azimuthal angles  $\phi_1$  and  $\phi_2$  are the angles between the scattering plane and the transverse hadron momenta  $\mathbf{p}_{T1(2)}$  around the thrust axis  $\hat{n}$ . The polar angle  $\theta_{th}$  is the angle between  $\hat{n}$  and the beam axis. Note that the difference between  $\mathbf{p}_{T1(2)}$  and  $\mathbf{P}_{hT}$  is that the latter is calculated with respect to the  $q\bar{q}$  axis.

$$F^{[n]}(z_i) \equiv \int d|\mathbf{k}_T|^2 \left[ \frac{|\mathbf{k}_T|}{M_i} \right]^n F(z_i, |\mathbf{k}_T|^2), \quad (3)$$

with  $n = 0, 1$ , and  $|\mathbf{k}_T|$  the transverse momentum of the quarks with respect to the hadrons they fragment into, which, in this frame, is related to the measurement of the transverse momenta of the two hadrons with respect to the thrust axis.

Despite the simple form of the  $R_{12}$  normalized rate, which involves only the product of moments of FFs, the RF12 frame comes with several downsides, among others of having to rely on Monte Carlo (MC) simulations when using the thrust axis as a proxy for the leading-order  $q\bar{q}$  axis. An alternative frame is the analogue of the Gottfried-Jackson frame [23,24] which uses the momentum of one hadron as a reference axis, and defines a single angle  $\phi_0$  between the plane containing the two hadron momenta and the plane defined by the beam and the reference axis. We refer to this frame as RF0 [11,12]. The corresponding normalized yield in the  $e^+e^-$  c.m. system is [23]

$$R_0(2\phi_0) = 1 + \frac{\sin^2\theta_2}{1 + \cos^2\theta_2} \cos 2\phi_0 \cdot \frac{\mathcal{F}[(2\hat{\mathbf{h}} \cdot \mathbf{k}_T \hat{\mathbf{h}} \cdot \mathbf{p}_T - \mathbf{k}_T \cdot \mathbf{p}_T) H_1^\perp \bar{H}_1^\perp]}{(M_1 M_2) \mathcal{F}[D_1 \bar{D}_1]}, \quad (4)$$

where  $\theta_2$  is the angle between the hadron used as reference and the beam axis  $\hat{\mathbf{h}}$  is the unit vector in the direction of the transverse momentum of the first hadron relative to the axis defined by the second hadron, and  $\mathcal{F}$  is used to denote the convolution integral

$$\mathcal{F}[X\bar{X}] \equiv \sum_q e_q^2 \int d^2\mathbf{k}_T d^2\mathbf{p}_T \delta^2(\mathbf{p}_T + \mathbf{k}_T - \mathbf{q}_T) \times X^q(z_1, z_1^2 \mathbf{k}_T^2) \bar{X}^q(z_2, z_2^2 \mathbf{p}_T^2), \quad (5)$$

with  $\mathbf{k}_T$ ,  $\mathbf{p}_T$ , and  $\mathbf{q}_T$  the transverse momentum of the fragmenting quark, antiquark, and virtual photon from  $e^+e^-$  annihilation, respectively, in the frame where the two hadrons are collinear, and  $X(\bar{X}) \equiv D_1(\bar{D}_1)$  or  $H_1^\perp(\bar{H}_1^\perp)$ . In this frame, specific assumptions on the  $\mathbf{k}_T$  dependence of the involved functions are necessary to explicitly evaluate the convolution integrals.

For this analysis we use a data sample of 468 fb $^{-1}$  [25] collected at the c.m. energy  $\sqrt{s} \approx 10.6$  GeV with the *BABAR* detector [26,27] at the SLAC National Accelerator Laboratory. We use tracks reconstructed in the silicon vertex detector and in the drift chamber (DCH) and identified as pions or kaons in the DCH and in the Cherenkov ring imaging detector (DIRC). Detailed MC simulation is used to study detector effects and to estimate contribution from various background sources. Hadronic

events are generated using the *Jetset* [28] package and undergo a full detector simulation based on *GEANT4* [29].

We make a tight selection of hadronic events in order to minimize biases due to detector acceptance and hard initial-state photon radiation (ISR), as they can introduce fake azimuthal modulations. Furthermore, final-state gluon ( $q\bar{q}g$ ) radiation also leads to angular asymmetries to be taken into account [23]. Requiring at least three charged tracks consistent with the  $e^+e^-$  primary vertex and a total visible energy of the event in the laboratory frame  $E_{\text{tot}} > 11$  GeV, we reject  $e^+e^- \rightarrow \tau^+\tau^-$  and two-photon backgrounds, as well as ISR ( $q\bar{q}g$ ) events with the photon (one jet) along the beam line. About 10% of ISR photons are within our detector acceptance, and we reject events with a photon candidate with energy above 2 GeV. We require an event thrust value  $T > 0.8$  to suppress  $q\bar{q}g$  and  $B\bar{B}$  events, and  $|\cos\theta_{\text{th}}| < 0.6$  so that most tracks are within the detector acceptance.

We assign randomly the positive direction of the thrust axis, and divide each event into two hemispheres by the plane perpendicular to it. To ensure tracks are assigned to the correct hemispheres, we require them to be within a  $45^\circ$  angle of the thrust axis and to have  $z > 0.15$ . A “tight” identification algorithm is used to identify kaons (pions), which is about 80% (90%) efficient and has misidentification rates below 10% (5%). We select those pions and kaons that lie within the DIRC acceptance region with a polar angle in laboratory frame  $0.45 \text{ rad} < \theta_{\text{lab}} < 2.46 \text{ rad}$ . To minimize backgrounds, such as  $e^+e^- \rightarrow \mu^+\mu^-\gamma$  followed by photon conversion, we require  $z < 0.9$ .

We construct all the possible pairs of selected tracks reconstructed in opposite thrust hemispheres, and we calculate the corresponding azimuthal angles  $\phi_1$ ,  $\phi_2$ , and  $\phi_0$  in the respective reference frames. In this way, we identify three different samples of hadron pairs:  $KK$ ,  $K\pi$ , and  $\pi\pi$ . To reduce low-energy gluon radiation and the contribution due to wrong hemispheres assignment, we require  $Q_t < 3.5 \text{ GeV}/c$ , where  $Q_t$  is the transverse momentum of the virtual photon from  $e^+e^-$  annihilation in the frame where the two hadrons are collinear [23].

The analysis is performed in intervals of hadron fractional energies with the following boundaries: 0.15, 0.2, 0.3, 0.5, 0.9, for a total of 16 two-dimensional  $(z_1, z_2)$  intervals.

For each of the three samples, we evaluate the normalized yield distributions  $R_{12}$  and  $R_0$  for unlike ( $U$ ), like ( $L$ ), and any charge combination ( $C$ ) of hadron pairs as a function of  $\phi_1 + \phi_2$  and  $2\phi_0$ , as shown in the left plot of Fig. 2 for  $KK$  pairs, for example. These combinations of charged hadrons contain different contributions of favored and disfavored FFs, where a favored (disfavored) process refers to the production of a hadron for which one (none) of the valence quarks is of the same kind as the fragmenting quark. In particular, by selecting  $KK$  pairs, we are able to study the favored contribution  $H_s^{\perp\text{fav}}$  of the strange quark, not accessible when considering  $\pi\pi$  pairs only.




FIG. 2 (color online). Distributions of normalized yields (left plot) for unlike ( $U$ ), like ( $L$ ), and any charge combination ( $C$ ) of  $KK$  pairs, and their double ratios (right plot) in RF12.

The normalized distributions can be parametrized with a cosine function:  $R_\alpha^i = b_\alpha + a_\alpha^i \cos\beta_\alpha$ , where  $\alpha = 0, 12$  indicates the reference frames,  $i = U, L, C$  the charge combination of hadron pairs, and  $\beta_{12(0)} = \phi_{12}(2\phi_0)$ .

The  $R_\alpha^i$  distributions are strongly affected by instrumental effects. In order to reduce the impact of the detector acceptance, as well as any remaining effect from gluon bremsstrahlung [23], we construct two double ratios (DRs) of normalized distributions,  $R_\alpha^U/R_\alpha^L$  and  $R_\alpha^U/R_\alpha^C$ . The two ratios give access to the same physical quantities as the independent  $R_\alpha^i$ , that is the favored and disfavored FFs, but in different combinations. We report the results for both kinds of DRs, which are strongly correlated since they are obtained by using the same data set. These are shown in the right plot of Fig. 2 for  $KK$  pairs in RF12. At first order, the double ratios are still parametrized by a function that is linear in the cosine of the corresponding combination of azimuthal angles:

$$R_\alpha^{ij} = \frac{R_\alpha^i}{R_\alpha^j} \simeq B_\alpha^{ij} + A_\alpha^{ij} \cdot \cos\beta_\alpha, \quad (6)$$

with  $B$  and  $A$  free parameters, and  $i, j = U, L, C$ . The constant term  $B$  must be consistent with unity, while  $A$  contains the information about the favored and disfavored Collins FFs.

We fit the binned  $R_\alpha^{ij}$  distributions independently for  $KK$ ,  $K\pi$ , and  $\pi\pi$  hadron pairs. Using the MC sample, we evaluate the  $K/\pi$  (mis)identification probabilities for the 16  $(z_1, z_2)$  intervals in each of the three samples. For example, the probability  $f_{KK}^{KK}$  that a true  $KK$  pair is reconstructed as  $KK$  pair is about 90% on average, slightly decreasing at higher momenta, while the probability  $f_{K\pi}^{KK}$  that a true  $K\pi$  pair is identified as  $KK$  is about 10%, and  $f_{\pi\pi}^{KK}$  is negligible.

The presence of background processes could introduce azimuthal modulations not related to the Collins effect, and modifies the measured asymmetry as follows:

$$A_{KK}^{\text{meas}} = F_{uds}^{KK} \cdot \left( \sum_{nm} f_{nm}^{KK} \cdot A_{nm} \right) + \sum_i F_i^{KK} \left( \sum_{nm} f_{nm}^{(KK)i} \cdot A_{nm}^i \right), \quad (7)$$

with  $nm = KK, K\pi, \pi\pi$ , and  $i = c\bar{c}, B\bar{B}, \tau^+\tau^-$ . In Eq. (7),  $A_{nm}$  are the true Collins asymmetries produced from the fragmentation of light quarks in the three samples,  $A_{nm}^i$  is the  $i$ th background asymmetry contribution, and  $F_{uds(i)}^{KK}$  are the fractions of reconstructed kaon pairs coming from  $uds$  and background events, calculated from the respective MC samples. By construction,  $\sum_i F_i + F_{uds} = 1$ . A similar expression holds for  $K\pi$  and  $\pi\pi$  samples.

Previous studies [11] show that  $e^+e^- \rightarrow B\bar{B}$  and  $\tau^+\tau^-$  events have negligible  $A_{nm}^i$ ,  $F_{B\bar{B}} < 2\%$ , and  $F_{\tau^+\tau^-}$  significantly different from zero only for the  $\pi\pi$  sample at high  $z$  values. Since  $F_{c\bar{c}}$  can be as large as 30%, and  $A_{c\bar{c}}$  are unknown, we determine  $A_{nm}^{\text{meas}}$  in Eq. (7) from samples enhanced in  $c\bar{c}$  by requiring the reconstruction of at least one  $D^{*\pm}$  meson from the decay  $D^{*\pm} \rightarrow D^0\pi^\pm$ , with the  $D^0$  candidate reconstructed in the following four Cabibbo-favored decay modes:  $K^-\pi^+$ ,  $K^-\pi^+\pi^-\pi^+$ ,  $K_s^0\pi^+\pi^-$ , and  $K^-\pi^+\pi^0$ . These modes are assumed to provide a representative sample of  $\pi\pi$ ,  $K\pi$ , and  $KK$  pairs to be used in the correction, an assumption that is strengthened by the observation that the background asymmetries for those modes were found to be consistent. We solve the system of equations for  $A_{KK}^{\text{meas}}$ ,  $A_{K\pi}^{\text{meas}}$ ,  $A_{\pi\pi}^{\text{meas}}$ , for the standard and charm-enhanced samples, and we extract simultaneously the Collins asymmetries  $A_{KK}$ ,  $A_{K\pi}$ , and  $A_{\pi\pi}$ , corrected for the contributions of the background and  $K/\pi$  (mis) identification. The dominant uncertainties related to this procedure come from the limited statistics of the  $D^*$ -enhanced sample and from the fractions  $F_i$ . The uncertainties on the fractions are evaluated by data-MC comparison and amount to a few percent. All these uncertainties are therefore included in the statistical error of the asymmetries extracted from the system of Eq. (7).

We test the DR method on the MC sample. Spin effects are not simulated in MC, and so the DR distributions should be uniform. However, when fitting the distributions for reconstructed  $KK$  pairs with Eq. (6), we measure a cosine term in the full sample of  $0.004 \pm 0.001$  and  $0.007 \pm 0.001$  in the RF12 and RF0 frames, respectively, indicating a bias. Smaller values are obtained for  $K\pi$  and  $\pi\pi$  pairs [30]. Studies performed on the MC samples, both at generation level and after full simulation, demonstrate that the main source of this bias is due to the emission of ISR, which boosts the hadronic system and distorts the angular distribution of the final state particles, resulting in azimuthal modulations not related to the Collins effect. This effect is more pronounced for  $KK$  pairs due to the lower multiplicity with respect to the other two combinations of hadrons. Assuming the bias, which is everywhere

smaller than the asymmetries measured in the data sample in each bin, is additive, we subtract it from the background-corrected asymmetry.

Using the  $uds$  MC sample, or light quark  $e^+e^- \rightarrow q\bar{q}$  MC events, we study the difference between measured and true azimuthal asymmetries. The asymmetry is introduced into the simulation by reweighting the events according to the distribution  $1 \pm a \cdot \cos \phi_a^{\text{gen}}$ , where we use different values of  $a$  ranging from 0 to 8% with positive (negative) sign for  $U$  ( $L$  and  $C$ ) hadron pairs, and  $\phi_a^{\text{gen}}$  are the azimuthal angles combinations calculated with respect to the true  $q\bar{q}$  axis in RF12, or the generated hadron momentum in RF0. The reconstructed asymmetries in RF12 are systematically underestimated for the three samples of hadron pairs, as expected since we use the thrust axis instead of the  $q\bar{q}$  axis, while they are consistent with the simulated ones in RF0, where only particle identification and tracking reconstruction effects could introduce possible dilution. Since we measure the same dilution for  $KK$ ,  $K\pi$ , and  $\pi\pi$  samples, the asymmetry is corrected by rescaling  $A_{KK}$ ,  $A_{K\pi}$ , and  $A_{\pi\pi}$  using the same correction factor, which ranges from 1.3 to 2.3 increasing with  $z$ , as shown in Fig. 3. No corrections are needed for the asymmetries measured in RF0. The uncertainties on the correction factors are assigned as systematic contributions.

All systematic effects, if not otherwise specified, are evaluated for each bin of  $z$ . The main contribution comes from the MC bias. We compare the bias results from the nominal selection, with those obtained by requiring different cuts on  $E_{\text{tot}}$ , and/or by changing the detector acceptance region for the hadrons. The largest variation of the bias is combined in quadrature with the MC statistical error and




FIG. 3 (color online). Correction factors for the dilution of the asymmetry due to the difference between the thrust and the  $q\bar{q}$  axis. The open (full) markers, triangles and circles, refer to the  $U/L$  and  $U/C$  double ratios in the RF12 (RF0) frame, respectively. The 16  $(z_1, z_2)$  bins are shown on the  $x$  axis: in each interval between the dashed lines,  $z_1$  is chosen in the following ranges: [0.15, 0.2], [0.2, 0.3], [0.3, 0.5], and [0.5, 0.9], while within each interval the points correspond to the four bins in  $z_2$ .

taken as systematic uncertainty. The effects due to the particle identification are evaluated using tighter and looser selection criteria. The largest deviations with respect to the nominal selection are taken as systematic uncertainties: the average relative uncertainties are around 10%, 7%, and 5% for the  $KK$ ,  $K\pi$ , and  $\pi\pi$  pairs. Fitting the azimuthal distributions using different bin sizes, we determine relative systematic uncertainties, which are not larger than 5%, 1.9%, and 1% for the three samples. The systematic uncertainty due to the  $E_{\text{tot}}$  cut is obtained by comparing the measured asymmetries with those obtained with the looser selection  $E_{\text{tot}} > 10$  GeV. The average systematic contribution is around 10% for the three samples in both reference frames. We use different fitting functions with additional higher harmonic terms. No significant changes in the value of the cosine moments with respect to the standard fits are found. As a cross-check of the double ratio method we fit the difference of  $R^i$  distributions, and we compare the two results. The difference between the two procedures is negligible for  $K\pi$  and  $\pi\pi$  pairs, while it reaches 1% and 3% for kaon pairs in RF12 and RF0, respectively. All the other systematic contributions are negligible [11].

The Collins asymmetries measured for the 16 two-dimensional  $(z_1, z_2)$  bins, for reconstructed  $KK$ ,  $K\pi$ , and  $\pi\pi$  hadron pairs, are shown in Fig. 4 for RF12 and RF0, and are summarized in tables reported in the Supplemental Material [30]. The asymmetries are corrected for the background contributions and  $K/\pi$  contamination following Eq. (7), the MC bias is subtracted, and the corrections due to the dilution effects are applied. The total systematic uncertainties are obtained by adding in quadrature the individual contributions, and are represented by the bands around the data points.

An increasing asymmetry with increasing hadron energies is visible for the  $U/L$  double ratio in both reference frames. The largest effects, but with less precision, are observed for  $KK$  pairs, for which  $A_{12}^{UL}$  is consistent with zero at low  $z$ , and reaches 22% in the last  $z$  bin, while somewhat smaller values are seen for  $\pi\pi$  and  $K\pi$  pairs. In particular, at low  $(z_1, z_2)$  bins  $A^{UL}$  for  $\pi\pi$  pairs is nonzero, in agreement with the behavior observed in [11]. The small differences between the two data sets are due to the different kinematic region selected after the cut on  $\cos \theta_{th}$ . The  $A^{UC}$  asymmetry is smaller than  $A^{UL}$  in all cases, and, for the  $KK$  pairs, the rise of the asymmetry with the hadron energies is not evident. We also note that the asymmetries for the  $KK$  pairs are larger than the others when the  $U/L$  ratio is considered, while they are at the same level, or lower, when they are extracted from the  $U/C$  ratio.

In summary, we have studied for the first time in  $e^+e^-$  annihilation the Collins asymmetry for inclusive production of  $KK$  and  $K\pi$  pairs as a function of  $(z_1, z_2)$  in two distinct reference frames. We measure the azimuthal modulation of the double ratios  $U/L$  and  $U/C$ , which are sensitive to the favored and disfavored Collins FFs for light quarks. We simultaneously extract also the Collins asymmetries for  $\pi\pi$  pairs, which are found to be in agreement with those obtained in previous studies [11,13]. The results reported in this paper and those obtained from SIDIS experiments can be used in a global analysis to extract the favored contribution of the strange quark, and to improve the knowledge on the  $u$  and  $d$  fragmentation processes [14–16].

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II2 colleagues, and for the




FIG. 4 (color online). Comparison of  $U/L$  (top) and  $U/C$  (bottom) Collins asymmetries in RF12 (left) and RF0 (right) for  $KK$ ,  $K\pi$ , and  $\pi\pi$  pairs. The statistical and systematic uncertainties are represented by the bars and the bands around the points, respectively. The 16  $(z_1, z_2)$  bins are shown on the  $x$  axis: in each interval between the dashed lines,  $z_1$  is chosen in the following ranges: [0.15, 0.2], [0.2, 0.3], [0.3, 0.5], and [0.5, 0.9], while within each interval the points correspond to the four bins in  $z_2$ .

substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (U.S.), NSERC (Canada), CEA and CNRS-IN2P3 (France), BMBF and

DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MES (Russia), MEC (Spain), and STFC (U.K.). Individuals have received support from the Marie Curie EIF (European Union) and the A.P. Sloan Foundation.

- 
- [1] J. C. Collins, *Nucl. Phys.* **B396**, 161 (1993).
  - [2] J. C. Collins and D. E. Soper, *Nucl. Phys.* **B193**, 381 (1981).
  - [3] J. Collins, D. E. Soper, and G. Sterman, *Nucl. Phys.* **B250**, 199 (1985).
  - [4] J. C. Collins and A. Metz, *Phys. Rev. Lett.* **93**, 252001 (2004).
  - [5] P. Sun and F. Yuan, *Phys. Rev. D* **88**, 114012 (2013).
  - [6] A. Airapetian *et al.* (HERMES Collaboration), *Phys. Rev. Lett.* **94**, 012002 (2005).
  - [7] A. Airapetian *et al.* (HERMES Collaboration), *Phys. Lett. B* **693**, 11 (2010).
  - [8] C. Adolph *et al.* (COMPASS Collaboration), *Phys. Lett. B* **717**, 376 (2012).
  - [9] M. Alekseev *et al.* (COMPASS Collaboration), *Phys. Lett. B* **673**, 127 (2009).
  - [10] A. Airapetian *et al.* (HERMES Collaboration), *Phys. Rev. D* **87**, 012010 (2013).
  - [11] J. Lees *et al.* (*BABAR* Collaboration), *Phys. Rev. D* **90**, 052003 (2014).
  - [12] R. Seidl, G. Perdekamp *et al.* (Belle Collaboration), *Phys. Rev. D* **78**, 032011 (2008).
  - [13] R. Seidl, G. Perdekamp *et al.* (Belle Collaboration), *Phys. Rev. D* **86**, 039905(E) (2012).
  - [14] A. Bacchetta, L. P. Gamberg, G. R. Goldstein, and A. Mukherjee, *Phys. Lett. B* **659**, 234 (2008).
  - [15] M. Anselmino, M. Boglione, U. D'Alesio, A. Kotzinian, F. Murgia, A. Prokudin, and C. Türk, *Phys. Rev. D* **75**, 054032 (2007).
  - [16] M. Anselmino, M. Boglione, U. D'Alesio, S. Melis, F. Murgia, and A. Prokudin, *Phys. Rev. D* **87**, 094019 (2013).
  - [17] R. L. Jaffe and X. Ji, *Phys. Rev. Lett.* **67**, 552 (1991).
  - [18] J. Soffer, *Phys. Rev. Lett.* **74**, 1292 (1995).
  - [19] A. Martin, F. Bradamante, and V. Barone, *Phys. Rev. D* **91**, 014034 (2015).
  - [20] A. Bacchetta, U. D'Alesio, M. Diehl, and C. A. Miller, *Phys. Rev. D* **70**, 117504 (2004).
  - [21] E. Farhi, *Phys. Rev. Lett.* **39**, 1587 (1977).
  - [22] S. Brandt, C. Peyrou, R. Sosnowski, and A. Wroblewski, *Phys. Lett.* **12**, 57 (1964).
  - [23] D. Boer, *Nucl. Phys.* **B806**, 23 (2009).
  - [24] D. Boer, R. Jakob, and P. Mulders, *Nucl. Phys.* **B504**, 345 (1997).
  - [25] J. P. Lees *et al.* (*BABAR* Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **726**, 203 (2013).
  - [26] B. Aubert *et al.* (*BABAR* Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **479**, 1 (2002).
  - [27] B. Aubert *et al.* (*BABAR* Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **729**, 615 (2013).
  - [28] T. Sjöstrand, *arXiv:hep-ph/9508391*.
  - [29] S. Agostinelli *et al.* (GEANT4 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
  - [30] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevD.92.111101> for tables and figures.

*Collins Asymmetries in Inclusive Charged  $KK$  and  $K\pi$  Pairs  
Produced in  $e^+e^-$  Annihilation*

©2015 American Physical Society

**Citation:**

Lees, J. P., V. Poireau, V. Tisserand, E. Grauges, et al. 2015. "Collins asymmetries in inclusive charged  $KK$  and  $K\pi$  pairs produced in  $e^+e^-$  annihilation." *Physical Review D - Particles, Fields, Gravitation and Cosmology* 92(11), doi:10.1103/PhysRevD.92.111101.

*This document is being made freely available by the Eugene McDermott Library of The University of Texas at Dallas with permission from the copyright owner. All rights are reserved under United States copyright law unless specified otherwise.*