

School of Natural Sciences and Mathematics

***Dark Photon Search in the Mass
Range between 1.5 and 3.4 GeV/c²***

UT Dallas Author(s):

Xinchou Lou

Rights:

CC BY 4.0 (Attribution)

©2017 The Authors

Citation:

Ablikim, M., M. N. Achasov, X. C. Ai, O. Albayrak, et al. 2017. "Dark photon search in the mass range between 1.5 and 3.4 GeV/c²." Physics Letters B 774, doi:10.1016/j.physletb.2017.09.067

This document is being made freely available by the Eugene McDermott Library of the University of Texas at Dallas with permission of the copyright owner. All rights are reserved under United States copyright law unless specified otherwise.

Dark photon search in the mass range between 1.5 and 3.4 GeV/c²

BESIII Collaboration

M. Ablikim^a, M.N. Achasov^{i,6}, X.C. Ai^a, O. Albayrak^e, M. Albrecht^d, D.J. Ambrose^{av}, A. Amoroso^{az,bb}, F.F. An^a, Q. An^{aw,1}, J.Z. Bai^a, R. Baldini Ferroli^t, Y. Ban^{ag}, D.W. Bennett^s, J.V. Bennett^e, M. Bertani^t, D. Bettoni^v, J.M. Bian^{au}, F. Bianchi^{az,bb}, E. Boger^{y,4}, I. Boyko^y, R.A. Briere^e, H. Cai^{bd}, X. Cai^{a,1}, O. Cakir^{ap,2}, A. Calcaterra^t, G.F. Cao^a, S.A. Cetin^{aq}, J.F. Chang^{a,1}, G. Chelkov^{y,4,5}, G. Chen^a, H.S. Chen^a, H.Y. Chen^b, J.C. Chen^a, M.L. Chen^{a,1}, S.J. Chen^{ae}, X. Chen^{a,1}, X.R. Chen^{ab}, Y.B. Chen^{a,1}, H.P. Cheng^q, X.K. Chu^{ag}, G. Cibinetto^v, H.L. Dai^{a,1}, J.P. Dai^{aj}, A. Dbeyssiⁿ, D. Dedovich^y, Z.Y. Deng^a, A. Denig^x, I. Denysenko^y, M. Destefanis^{az,bb}, F. De Mori^{az,bb}, Y. Ding^{ac}, C. Dong^{af}, J. Dong^{a,1}, L.Y. Dong^a, M.Y. Dong^{a,1}, S.X. Du^{bf}, P.F. Duan^a, E.E. Eren^{aq}, J.Z. Fan^{ao}, J. Fang^{a,1}, S.S. Fang^a, X. Fang^{aw,1}, Y. Fang^a, L. Fava^{ba,bb}, F. Feldbauer^x, G. Felici^t, C.Q. Feng^{aw,1}, E. Fioravanti^v, M. Fritsch^{n,x}, C.D. Fu^a, Q. Gao^a, X.Y. Gao^b, Y. Gao^{ao}, Z. Gao^{aw,1}, I. Garzia^v, C. Geng^{aw,1}, K. Goetzen^j, W.X. Gong^{a,1}, W. Gradl^x, M. Greco^{az,bb}, M.H. Gu^{a,1}, Y.T. Gu^l, Y.H. Guan^a, A.Q. Guo^a, L.B. Guo^{ad}, Y. Guo^a, Y.P. Guo^x, Z. Haddadi^{aa}, A. Hafner^x, S. Han^{bd}, Y.L. Han^a, X.Q. Hao^o, F.A. Harris^{at}, K.L. He^a, Z.Y. He^{af}, T. Held^d, Y.K. Heng^{a,1}, Z.L. Hou^a, C. Hu^{ad}, H.M. Hu^a, J.F. Hu^{az,bb}, T. Hu^{a,1}, Y. Hu^a, G.M. Huang^f, G.S. Huang^{aw,1}, H.P. Huang^{bd}, J.S. Huang^o, X.T. Huang^{ai}, Y. Huang^{ae}, T. Hussain^{ay}, Q. Ji^a, Q.P. Ji^{af}, X.B. Ji^a, X.L. Ji^{a,1}, L.L. Jiang^a, L.W. Jiang^{bd}, X.S. Jiang^{a,1}, X.Y. Jiang^{af}, J.B. Jiao^{ai}, Z. Jiao^q, D.P. Jin^{a,1}, S. Jin^a, T. Johansson^{bc}, A. Julin^{au}, N. Kalantar-Nayestanaki^{aa}, X.L. Kang^a, X.S. Kang^{af}, M. Kavatsyuk^{aa}, B.C. Ke^e, P. Kiese^x, R. Kliemtⁿ, B. Kloss^x, O.B. Kolcu^{aq,9}, B. Kopf^d, M. Kornicer^{at}, W. Kuehn^z, A. Kupsc^{bc}, J.S. Lange^z, M. Lara^s, P. Larinⁿ, C. Leng^{bb}, C. Li^{bc}, C.H. Li^a, Cheng Li^{aw,1}, D.M. Li^{bf}, F. Li^{a,1}, G. Li^a, H.B. Li^a, J.C. Li^a, Jin Li^{ah}, K. Li^{ai}, K. Li^m, Lei Li^c, P.R. Li^{as}, T. Li^{ai}, W.D. Li^a, W.G. Li^a, X.L. Li^{ai}, X.M. Li^l, X.N. Li^{a,1}, X.Q. Li^{af}, Z.B. Li^{an}, H. Liang^{aw,1}, Y.F. Liang^{al}, Y.T. Liang^z, G.R. Liao^k, D.X. Linⁿ, B.J. Liu^a, C.X. Liu^a, F.H. Liu^{ak}, Fang Liu^a, Feng Liu^f, H.B. Liu^l, H.H. Liu^p, H.H. Liu^a, H.M. Liu^a, J. Liu^a, J.B. Liu^{aw,1}, J.P. Liu^{bd}, J.Y. Liu^a, K. Liu^{ao}, K.Y. Liu^{ac}, L.D. Liu^{ag}, P.L. Liu^{a,1}, Q. Liu^{as}, S.B. Liu^{aw,1}, X. Liu^{ab}, X.X. Liu^{as}, Y.B. Liu^{af}, Z.A. Liu^{a,1}, Zhiqiang Liu^a, Zhiqing Liu^x, H. Loehner^{aa}, X.C. Lou^{a,1,8}, H.J. Lu^q, J.G. Lu^{a,1}, R.Q. Lu^r, Y. Lu^a, Y.P. Lu^{a,1}, C.L. Luo^{ad}, M.X. Luo^{be}, T. Luo^{at}, X.L. Luo^{a,1}, M. Lv^a, X.R. Lyu^{as}, F.C. Ma^{ac}, H.L. Ma^a, L.L. Ma^{ai}, Q.M. Ma^a, T. Ma^a, X.N. Ma^{af}, X.Y. Ma^{a,1}, F.E. Maasⁿ, M. Maggiora^{az,bb}, Y.J. Mao^{ag}, Z.P. Mao^a, S. Marcello^{az,bb}, J.G. Messchendorp^{aa}, J. Min^{a,1}, T.J. Min^a, R.E. Mitchell^s, X.H. Mo^{a,1}, Y.J. Mo^f, C. Morales Moralesⁿ, K. Moriya^s, N.Yu. Muchnoi^{i,6}, H. Muramatsu^{au}, Y. Nefedov^y, F. Nerlingⁿ, I.B. Nikolaev^{i,6}, Z. Ning^{a,1}, S. Nisar^h, S.L. Niu^{a,1}, X.Y. Niu^a, S.L. Olsen^{ah}, Q. Ouyang^{a,1}, S. Pacetti^u, P. Patteri^t, M. Pelizaeus^d, H.P. Peng^{aw,1}, K. Peters^j, J. Pettersson^{bc}, J.L. Ping^{ad}, R.G. Ping^a, R. Poling^{au}, V. Prasad^a, Y.N. Pu^r, M. Qi^{ae}, S. Qian^{a,1}, C.F. Qiao^{as}, L.Q. Qin^{ai}, N. Qin^{bd}, X.S. Qin^a, Y. Qin^{ag}, Z.H. Qin^{a,1}, J.F. Qiu^a

E-mail address: guo@uni-mainz.de (Y.P. Guo).

<https://doi.org/10.1016/j.physletb.2017.09.067>0370-2693/© 2017 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

K.H. Rashid^{ay}, C.F. Redmer^x, H.L. Ren^r, M. Ripka^x, G. Rong^a, Ch. Rosnerⁿ, X.D. Ruan^l, V. Santoro^v, A. Sarantsev^{y,7}, M. Savrié^w, K. Schoenning^{bc}, S. Schumann^x, W. Shan^{ag}, M. Shao^{aw,1}, C.P. Shen^b, P.X. Shen^{af}, X.Y. Shen^a, H.Y. Sheng^a, W.M. Song^a, X.Y. Song^a, S. Sosio^{az,bb}, S. Spataro^{az,bb}, G.X. Sun^a, J.F. Sun^o, S.S. Sun^a, Y.J. Sun^{aw,1}, Y.Z. Sun^a, Z.J. Sun^{a,1}, Z.T. Sun^s, C.J. Tang^{al}, X. Tang^a, I. Tapan^{ar}, E.H. Thorndike^{av}, M. Tiemens^{aa}, M. Ullrich^z, I. Uman^{aq}, G.S. Varner^{at}, B. Wang^{af}, B.L. Wang^{as}, D. Wang^{ag}, D.Y. Wang^{ag}, K. Wang^{a,1}, L.L. Wang^a, L.S. Wang^a, M. Wang^{ai}, P. Wang^a, P.L. Wang^a, S.G. Wang^{ag}, W. Wang^{a,1}, X.F. Wang^{ao}, Y.D. Wangⁿ, Y.F. Wang^{a,1}, Y.Q. Wang^x, Z. Wang^{a,1}, Z.G. Wang^{a,1}, Z.H. Wang^{aw,1}, Z.Y. Wang^a, T. Weber^x, D.H. Wei^k, J.B. Wei^{ag}, P. Weidenkaff^x, S.P. Wen^a, U. Wiedner^d, M. Wolke^{bc}, L.H. Wu^a, Z. Wu^{a,1}, L.G. Xia^{ao}, Y. Xia^r, D. Xiao^a, H. Xiao^{ax}, Z.J. Xiao^{ad}, Y.G. Xie^{a,1}, Q.L. Xiu^{a,1}, G.F. Xu^a, L. Xu^a, Q.J. Xu^m, Q.N. Xu^{as}, X.P. Xu^{am}, L. Yan^{aw,1}, W.B. Yan^{aw,1}, W.C. Yan^{aw,1}, Y.H. Yan^r, H.J. Yang^{aj}, H.X. Yang^a, L. Yang^{bd}, Y. Yang^f, Y.X. Yang^k, H. Ye^a, M. Ye^{a,1}, M.H. Ye^g, J.H. Yin^a, B.X. Yu^{a,1}, C.X. Yu^{af}, H.W. Yu^{ag}, J.S. Yu^{ab}, C.Z. Yuan^a, W.L. Yuan^{ae}, Y. Yuan^a, A. Yuncu^{aq,3}, A.A. Zafar^{ay}, A. Zallo^t, Y. Zeng^r, B.X. Zhang^a, B.Y. Zhang^{a,1}, C. Zhang^{ae}, C.C. Zhang^a, D.H. Zhang^a, H.H. Zhang^{an}, H.Y. Zhang^{a,1}, J.J. Zhang^a, J.L. Zhang^a, J.Q. Zhang^a, J.W. Zhang^{a,1}, J.Y. Zhang^a, J.Z. Zhang^a, K. Zhang^a, L. Zhang^a, S.H. Zhang^a, X.Y. Zhang^{ai}, Y. Zhang^a, Y.N. Zhang^{as}, Y.H. Zhang^{a,1}, Y.T. Zhang^{aw,1}, Yu Zhang^{as}, Z.H. Zhang^f, Z.P. Zhang^{aw}, Z.Y. Zhang^{bd}, G. Zhao^a, J.W. Zhao^{a,1}, J.Y. Zhao^a, J.Z. Zhao^{a,1}, Lei Zhao^{aw,1}, Ling Zhao^a, M.G. Zhao^{af}, Q. Zhao^a, Q.W. Zhao^a, S.J. Zhao^{bf}, T.C. Zhao^a, Y.B. Zhao^{a,1}, Z.G. Zhao^{aw,1}, A. Zhemchugov^{y,4}, B. Zheng^{ax}, J.P. Zheng^{a,1}, W.J. Zheng^{ai}, Y.H. Zheng^{as}, B. Zhong^{ad}, L. Zhou^{a,1}, Li Zhou^{af}, X. Zhou^{bd}, X.K. Zhou^{aw,1}, X.R. Zhou^{aw,1}, X.Y. Zhou^a, K. Zhu^a, K.J. Zhu^{a,1}, S. Zhu^a, X.L. Zhu^{ao}, Y.C. Zhu^{aw,1}, Y.S. Zhu^a, Z.A. Zhu^a, J. Zhuang^{a,1}, L. Zotti^{az,bb}, B.S. Zou^a, J.H. Zou^a

^a Institute of High Energy Physics, Beijing 100049, People's Republic of China

^b Beihang University, Beijing 100191, People's Republic of China

^c Beijing Institute of Petrochemical Technology, Beijing 102617, People's Republic of China

^d Bochum Ruhr-University, D-44780 Bochum, Germany

^e Carnegie Mellon University, Pittsburgh, PA 15213, USA

^f Central China Normal University, Wuhan 430079, People's Republic of China

^g China Center of Advanced Science and Technology, Beijing 100190, People's Republic of China

^h COMSATS Institute of Information Technology, Lahore, Defence Road, Off Raiwind Road, 54000 Lahore, Pakistan

ⁱ G.I. Budker Institute of Nuclear Physics SB RAS (BINP), Novosibirsk 630090, Russia

^j GSI Helmholtz Centre for Heavy Ion Research GmbH, D-64291 Darmstadt, Germany

^k Guangxi Normal University, Guilin 541004, People's Republic of China

^l Guangxi University, Nanning 530004, People's Republic of China

^m Hangzhou Normal University, Hangzhou 310036, People's Republic of China

ⁿ Helmholtz Institute Mainz, Johann-Joachim-Becher-Weg 45, D-55099 Mainz, Germany

^o Henan Normal University, Xinxiang 453007, People's Republic of China

^p Henan University of Science and Technology, Luoyang 471003, People's Republic of China

^q Huangshan College, Huangshan 245000, People's Republic of China

^r Hunan University, Changsha 410082, People's Republic of China

^s Indiana University, Bloomington, IN 47405, USA

^t INFN Laboratori Nazionali di Frascati, I-00044, Frascati, Italy

^u INFN and University of Perugia, I-06100, Perugia, Italy

^v INFN Sezione di Ferrara, I-44122, Ferrara, Italy

^w University of Ferrara, I-44122, Ferrara, Italy

^x Johannes Gutenberg University of Mainz, Johann-Joachim-Becher-Weg 45, D-55099 Mainz, Germany

^y Joint Institute for Nuclear Research, 141980 Dubna, Moscow region, Russia

^z Justus Liebig University Giessen, II. Physikalisches Institut, Heinrich-Buff-Ring 16, D-35392 Giessen, Germany

^{aa} KVI-CART, University of Groningen, NL-9747 AA Groningen, The Netherlands

^{ab} Lanzhou University, Lanzhou 730000, People's Republic of China

^{ac} Liaoning University, Shenyang 110036, People's Republic of China

^{ad} Nanjing Normal University, Nanjing 210023, People's Republic of China

^{ae} Nanjing University, Nanjing 210093, People's Republic of China

^{af} Nankai University, Tianjin 300071, People's Republic of China

^{ag} Peking University, Beijing 100871, People's Republic of China

^{ah} Seoul National University, Seoul 151-747, Republic of Korea

^{ai} Shandong University, Jinan 250100, People's Republic of China

^{aj} Shanghai Jiao Tong University, Shanghai 200240, People's Republic of China

^{ak} Shanxi University, Taiyuan 030006, People's Republic of China

^{al} Sichuan University, Chengdu 610064, People's Republic of China

^{am} Soochow University, Suzhou 215006, People's Republic of China

^{an} Sun Yat-Sen University, Guangzhou 510275, People's Republic of China

^{ao} Tsinghua University, Beijing 100084, People's Republic of China

^{ap} Istanbul Aydin University, 34295 Sefakoy, Istanbul, Turkey

^{aq} Dogus University, 34722 Istanbul, Turkey^{ar} Uludag University, 16059 Bursa, Turkey^{as} University of Chinese Academy of Sciences, Beijing 100049, People's Republic of China^{at} University of Hawaii, Honolulu, HI 96822, USA^{au} University of Minnesota, Minneapolis, MN 55455, USA^{av} University of Rochester, Rochester, NY 14627, USA^{aw} University of Science and Technology of China, Hefei 230026, People's Republic of China^{ax} University of South China, Hengyang 421001, People's Republic of China^{ay} University of the Punjab, Lahore-54590, Pakistan^{az} University of Turin, I-10125, Turin, Italy^{ba} University of Eastern Piedmont, I-15121, Alessandria, Italy^{bb} INFN, I-10125, Turin, Italy^{bc} Uppsala University, Box 516, SE-75120 Uppsala, Sweden^{bd} Wuhan University, Wuhan 430072, People's Republic of China^{be} Zhejiang University, Hangzhou 310027, People's Republic of China^{bf} Zhengzhou University, Zhengzhou 450001, People's Republic of China

ARTICLE INFO

Article history:

Received 11 May 2017

Received in revised form 19 July 2017

Accepted 22 September 2017

Available online 28 September 2017

Editor: V. Metag

Keywords:

Dark photon search

Initial state radiation

BESIII

ABSTRACT

Using a data set of 2.93 fb^{-1} taken at a center-of-mass energy $\sqrt{s} = 3.773 \text{ GeV}$ with the BESIII detector at the BEPCII collider, we perform a search for an extra $U(1)$ gauge boson, also denoted as a dark photon. We examine the initial state radiation reactions $e^+e^- \rightarrow e^+e^- \gamma_{\text{ISR}}$ and $e^+e^- \rightarrow \mu^+\mu^- \gamma_{\text{ISR}}$ for this search, where the dark photon would appear as an enhancement in the invariant mass distribution of the leptonic pairs. We observe no obvious enhancement in the mass range between 1.5 and $3.4 \text{ GeV}/c^2$ and set a 90% confidence level upper limit on the mixing strength of the dark photon and the Standard Model photon. We obtain a competitive limit in the tested mass range.

© 2017 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

Several astrophysical anomalies, which cannot be easily understood in the context of the Standard Model (SM) of particle physics or astrophysics, have been discussed in relation to a dark, so far unobserved sector [1], which couples very weakly with SM particles. The most straightforward model consists of an extra $U(1)$ force carrier, also denoted as a dark photon, γ' , which couples to the SM via kinetic mixing [2]. It has been shown in Ref. [1] that the dark photon has to be relatively light, on the MeV/c^2 to GeV/c^2 mass scale, to explain the astrophysical observations. Furthermore, it was realized, that a dark photon of similar mass could also explain the presently observed deviation on the level of 3 to 4σ between the measurement and the SM prediction of $(g-2)_\mu$ [3]. These facts and the work by Bjorken and collaborators [4] triggered searches for the dark photon at particle accelerators in a world wide effort [5,6]. Different experimental setups can be used, like fixed-target (e.g. Refs. [7,8]), beam dump (e.g. Refs. [9,10]), or low-energy collider experiments (e.g. Refs. [11,12]). The mixing strength $\varepsilon = \alpha'/\alpha$, where α' is the coupling of the dark photon to the electromagnetic charge and α the fine structure constant, is constrained by previous measurements to be below approximately 10^{-2} [4].

In this letter we present a dark photon search, using 2.93 fb^{-1} [13] of data taken at $\sqrt{s} = 3.773 \text{ GeV}$ obtained with the Beijing Spectrometer III (BESIII). The measurement exploits the process of

initial state radiation (ISR), in which one of the beam particles radiates a photon. In this way, the available energy to produce final states is reduced, and the di-lepton invariant masses below the center-of-mass energy of the e^+e^- collider become available. The same method has been used by the BaBar experiment [11, 12], where a dark photon mass $m_{\gamma'}$ between 0.02 and $10.2 \text{ GeV}/c^2$ and ε values in the order of 10^{-3} – 10^{-4} have been excluded. We search for the processes $e^+e^- \rightarrow \gamma' \gamma_{\text{ISR}} \rightarrow l^+l^- \gamma_{\text{ISR}}$ ($l = \mu, e$) with leptonic invariant masses m_{l+l^-} between 1.5 and $3.4 \text{ GeV}/c^2$. The ISR QED processes $e^+e^- \rightarrow \mu^+\mu^- \gamma_{\text{ISR}}$ and $e^+e^- \rightarrow e^+e^- \gamma_{\text{ISR}}$ are irreducible background channels. However, the dark photon width is expected to be smaller than the resolution of the experiment [4] and, thus, a γ' signal would lead to a narrow structure at the mass of the dark photon in the m_{l+l^-} mass spectrum on top of the continuum QED background.

The BESIII detector is located at the double-ring e^+e^- Beijing Electron Positron Collider (BEPCII) [14]. The cylindrical BESIII detector covers 93% of the full solid angle. It consists of the following detector systems. (1) A Multilayer Drift Chamber (MDC) filled with a helium-gas mixture, composed of 43 layers, which provides a spatial resolution of $135 \mu\text{m}$ and a momentum resolution of 0.5% for charged tracks at $1 \text{ GeV}/c$ in a magnetic field of 1 T. (2) A Time-of-Flight system (TOF), built with 176 plastic scintillator counters in the barrel part, and 96 counters in the end caps. The time resolution in the barrel (end caps) is 80 ps (110 ps). For momenta up to $1 \text{ GeV}/c$, this provides a 2σ K/π separation. (3) A CsI(Tl) Electro-Magnetic Calorimeter (EMC) with an energy resolution of 2.5% in the barrel and 5% in the end caps at an energy of 1 GeV. (4) A Muon Counter (MUC) consisting of nine barrel and eight end-cap resistive plate chamber layers with a 2 cm position resolution.

For the simulation of ISR processes $e^+e^- \rightarrow \mu^+\mu^- \gamma_{\text{ISR}}$ and $\pi^+\pi^- \gamma_{\text{ISR}}$, the PHOKHARA event generator [15,16], which includes ISR and final state radiation (FSR) corrections up to next-to-leading order, is used. Bhabha scattering is simulated with BABAYAGA 3.5 [17]. Continuum Monte Carlo (MC) events, as well as the resonant $\psi(3770)$ decays to $D\bar{D}$, non- $D\bar{D}$, and the ISR

¹ Also at State Key Laboratory of Particle Detection and Electronics, Beijing 100049, Hefei 230026, People's Republic of China.

² Also at Ankara University, 06100 Tandogan, Ankara, Turkey.

³ Also at Bogazici University, 34342 Istanbul, Turkey.

⁴ Also at the Moscow Institute of Physics and Technology, Moscow 141700, Russia.

⁵ Also at the Functional Electronics Laboratory, Tomsk State University, Tomsk, 634050, Russia.

⁶ Also at the Novosibirsk State University, Novosibirsk, 630090, Russia.

⁷ Also at the NRC "Kurchatov Institute", PNPI, 188300, Gatchina, Russia.

⁸ Also at University of Texas at Dallas, Richardson, TX 75083, USA.

⁹ Currently at Istanbul Arel University, 34295 Istanbul, Turkey.




Fig. 1. Leptonic invariant mass distributions $m_{\mu^+\mu^-}$ and $m_{e^+e^-}$ after applying the selection requirements. Shown is data (points) and MC simulation (shaded area), which is scaled to the luminosity of the data set. The marked area around the J/ψ resonance is excluded in the analysis. The lower panel shows the ratio of fit curve and MC simulation (points) and the ratio of fit curve and MC simulation (histogram).

production of ψ' and J/ψ , are simulated with the KKMC generator [18]. All MC generators, which are the most appropriate choices for the processes studied, have been interfaced with the GEANT4-based [19,20] detector simulation.

The selection of $\mu^+\mu^-\gamma_{\text{ISR}}$ and $e^+e^-\gamma_{\text{ISR}}$ events is straightforward. We require the presence of two charged tracks in the MDC with net charge zero. The points of closest approach from the interaction point (IP) for these two tracks are required to be within a cylinder of 1 cm radius in the transverse direction and ± 10 cm of length along the beam axis. The polar angle with respect to the beam axis θ of the tracks is required to be in the fiducial volume of the MDC: $0.4 < \theta < \pi - 0.4$ radians. In order to suppress spiraling tracks, we require the transverse momentum p_t to be above 300 MeV/c for both tracks.

Muon particle identification is used [21]. The probabilities for being a muon $P(\mu)$ and being an electron $P(e)$ are calculated using information from MDC, TOF, EMC, and MUC. For both charged tracks, $P(\mu) > P(e)$ is required. To select electrons, the ratio of the measured energy in the EMC, E , to the momentum p obtained from the MDC is used. Both charged tracks must satisfy $E/p > 0.8$ c.

The radiator function [22], which describes the radiation of an ISR photon, is peaked at small θ values with respect to the beam axis. Different from BaBar, we use untagged ISR events, where the ISR photon is emitted at a small angle θ_γ and is not detected within the angular acceptance of the EMC, to increase statistics. A one constraint (1C) kinematic fit, applying energy and momentum conservation, is performed with the hypothesis $e^+e^- \rightarrow \mu^+\mu^-\gamma_{\text{ISR}}$ or $e^+e^- \rightarrow e^+e^-\gamma_{\text{ISR}}$, using as input the two selected charged track candidates, as well as the four momentum of the initial e^+e^- system. The constraint is the mass of a missing photon. The fit quality condition $\chi^2_{1C}/(\text{dof} = 1) < 20$ is applied in the $\mu^+\mu^-\gamma_{\text{ISR}}$ case, where dof is the degree of freedom. To suppress non-ISR background, the angle of the missing photon, θ_γ , predicted by the 1C kinematic fit, is required to be smaller than 0.1 radians or greater than $\pi - 0.1$ radians. We apply stronger requirements for the $e^+e^-\gamma_{\text{ISR}}$ final state, to provide a better suppression of the non-ISR background which is higher in the e^+e^- channel compared to the $\mu^+\mu^-$ channel. In this case, $\chi^2_{1C}/(\text{dof} = 1) < 5$, and $\theta_\gamma < 0.05$ radians, or $\theta_\gamma > \pi - 0.05$ radians.

Background in addition to the radiative QED processes $\mu^+\mu^-\gamma_{\text{ISR}}$ and $e^+e^-\gamma_{\text{ISR}}$, which is irreducible, is studied with MC simulations and is negligible for the $e^+e^-\gamma_{\text{ISR}}$ final state, and on the order of 3% for $\mu^+\mu^-$ invariant masses below 2 GeV/c² due to muon misidentification, and negligible above. This remaining back-

ground comes mostly from $\pi^+\pi^-\gamma_{\text{ISR}}$ events. We subtract their contribution using a MC sample, produced with the PHOKHARA generator. The subtraction of this background leads to a systematic uncertainty due to the generator precision smaller than 0.5%.

The $\mu^+\mu^-$ and e^+e^- invariant mass distributions, $m_{\mu^+\mu^-}$ and $m_{e^+e^-}$, which are shown separately in Fig. 1, are mainly dominated by the QED background but could contain the signal sitting on top of these irreducible events. For comparison with data, MC simulation, scaled to the luminosity of data, is shown, although it is not used in the search for the dark photon. In this analysis, the dark photon mass range $m_{\gamma'}$ between 1.5 and 3.4 GeV/c² is studied. Below 1.5 GeV/c² the $\pi^+\pi^-\gamma_{\text{ISR}}$ cross section with muon misidentification dominates the $m_{\mu^+\mu^-}$ spectrum. Above 3.4 GeV/c² the hadronic $q\bar{q}$ process can not be suppressed sufficiently by the χ^2_{1C} requirement. In order to search for narrow structures on top of the QED background, 4th order polynomial functions to describe the continuum QED are fitted to the data distributions shown in Fig. 1. The mass range around the narrow J/ψ resonance between 2.95 and 3.2 GeV/c² is excluded.

The differences between the $\mu^+\mu^-\gamma_{\text{ISR}}$ and $e^+e^-\gamma_{\text{ISR}}$ event yields and their respective 4th order polynomials are added. The combined differences are represented by the black dots in Fig. 2. A dark photon candidate would appear as a peak in this plot. The observed statistical significances are less than 3σ everywhere in the explored region. The significance in each invariant mass bin is defined as the combined differences between data and the 4th order polynomials, divided by the combined statistical errors of both final states. In conclusion, we observe no dark photon signal for $1.5 \text{ GeV/c}^2 < m_{\gamma'} < 3.4 \text{ GeV/c}^2$, where $m_{\gamma'}$ is equal to the leptonic invariant mass $m_{l^+l^-}$. The exclusion limit at the 90% confidence level is determined with a profile likelihood approach [23]. Also shown in Fig. 2 as a function of $m_{l^+l^-}$ is the bin-by-bin calculated exclusion limit, including the systematic uncertainties as explained below.

To calculate the exclusion limit on the mixing parameter ε^2 , the formula from Ref. [4] is used

$$\frac{\sigma_i(e^+e^- \rightarrow \gamma' \gamma_{\text{ISR}} \rightarrow l^+l^- \gamma_{\text{ISR}})}{\sigma_i(e^+e^- \rightarrow \gamma^* \gamma_{\text{ISR}} \rightarrow l^+l^- \gamma_{\text{ISR}})} = \frac{N_i^{\text{up}}(e^+e^- \rightarrow \gamma' \gamma_{\text{ISR}} \rightarrow l^+l^- \gamma_{\text{ISR}})}{N_i^{\text{B}}(e^+e^- \rightarrow \gamma^* \gamma_{\text{ISR}} \rightarrow l^+l^- \gamma_{\text{ISR}})} \cdot \frac{1}{\varepsilon} = \frac{3\pi \cdot \varepsilon^2 \cdot m_{\gamma'}}{2N_f^{l^+l^-} \alpha \cdot \delta_m^{l^+l^-}}, \quad (1)$$




Fig. 2. The sum of the differences between the $\mu^+\mu^-\gamma_{\text{ISR}}$ and $e^+e^-\gamma_{\text{ISR}}$ event yields and their respective 4th order polynomials (dots with error bars). The solid histogram represents the exclusion limit with the 90% confidence, calculated with a profile likelihood approach and including the systematic uncertainty. The region around the J/ψ resonance between 2.95 and 3.2 GeV/c^2 is excluded.

where i represents the i -th mass bin, α is the electromagnetic fine structure constant, $m_{\gamma'}$ the dark photon mass, γ^* the SM photon, and $\delta_m^{l+l^-}$ ($l = \mu, e$) the bin width of the lepton pair invariant mass spectrum, $10 \text{ MeV}/c^2$. The mass resolution of the lepton pairs determined with MC for e^+e^- and $\mu^+\mu^-$ is between 5 and $12 \text{ MeV}/c^2$. The cross section ratio upper limit in Eq. (1) is determined from the exclusion upper limit (N^{up}) corrected by the efficiency loss (ϵ) due to the bin width divided by the number of $\mu^+\mu^-\gamma_{\text{ISR}}$ and $e^+e^-\gamma_{\text{ISR}}$ events (N^{B}) corrected as described below. The efficiency loss caused by the incompleteness of signal events in one bin is calculated with $\int_{-5}^5 \text{MeV}/c^2 G(0, \sigma) dm / \int_{-\infty}^{\infty} G(0, \sigma) dm$, where $G(0, \sigma)$ is the Gaussian function used to describe the mass resolution.

The QED cross section $\sigma_i(e^+e^- \rightarrow \gamma^* \gamma_{\text{ISR}} \rightarrow l^+l^- \gamma_{\text{ISR}})$ must only take into account annihilation processes of the initial e^+e^- beam particles, where a dark photon could be produced. Thus, the event yield of the $e^+e^- \gamma$ final state has to be corrected due to the existence of SM Bhabha scattering. This correction is obtained in bins of $m_{e^+e^-}$ by dividing the e^+e^- annihilation events only by the sum of events of the annihilation and Bhabha scattering processes. The first is generated with the PHOKHARA event generator by generating the $\mu^+\mu^-\gamma$ final state and replacing the muon mass with the electron mass. The latter is generated with the BABAYAGA@NLO

generator [24]. The correction factor varies between 2% and 8% depending on $m_{e^+e^-}$.

The number of final states for the dark photon $N_f^{l+l^-}$ includes the phase space above the l^+l^- production threshold of the leptons $l = \mu, e$, and is given by $N_f^{l+l^-} = \Gamma_{\text{tot}}/\Gamma_{ll}$ [25], where $\Gamma_{ll} \equiv \Gamma(\gamma' \rightarrow l^+l^-)$ is the leptonic γ' width and Γ_{tot} is the total γ' width. These widths are taken from Ref. [25]

$$\Gamma_{ll} = \frac{\alpha \varepsilon^2}{3m_{\gamma'}^2} (m_{\gamma'}^2 + 2m_l^2) \sqrt{m_{\gamma'}^2 - 4m_l^2} \quad (2)$$

$$\Gamma_{\text{tot}} = \Gamma_{ee} + \Gamma_{\mu\mu} \cdot (1 + R(\sqrt{s})), \quad (3)$$

where $\Gamma_{ee} \equiv \Gamma(\gamma' \rightarrow e^+e^-)$, $\Gamma_{\mu\mu} \equiv \Gamma(\gamma' \rightarrow \mu^+\mu^-)$, and $R(\sqrt{s})$ is the total hadronic cross section R value [26] as a function of \sqrt{s} .

The systematic uncertainties are included in the calculation of the exclusion limit. The main source is the uncertainty of the R value taken from Ref. [26], which enters the calculation of the $N_f^{l+l^-}$ and leads to a mass dependent systematic uncertainty between 3.0 and 6.0%. Other sources are background subtraction as described above ($<0.5\%$), the fitting error of the polynomial fit to data ($<1\%$), the Bhabha scattering correction factor using the PHOKHARA and BABAYAGA@NLO event generator ($<1\%$), and data-MC differences of the leptonic mass resolution. To quantify the latter one, we study the data-MC resolution difference of the J/ψ resonance for the $\mu^+\mu^-$ and e^+e^- decays, separately. The resonance is fitted with a double Gaussian function in data and MC simulation, and the width difference is $(3.7 \pm 1.8)\%$ for $\mu^+\mu^-$ and $(0.7 \pm 5.3)\%$ for e^+e^- . The differences are taken into consideration in the calculations, and the uncertainty in the differences (1%) is taken as the systematic uncertainty of the data-MC differences. The mass dependent total systematic uncertainty, which varies from 3.5 to 6.5% depending on mass, is used bin-by-bin in the upper limit.

The final result, the mixing strength ε as a function of the dark photon mass, is shown in Fig. 3, including the systematic uncertainties. It provides a comparable upper limit to BaBar [11,12] in the studied $m_{\gamma'}$ mass range. Also shown are the exclusion limits from KLOE [27–30], WASA-at-COSY [31], HADES [32], PHENIX [33], A1 at MAMI [7,8], NA48/2 [34], APEX [35], and the beam-dump experiments E774 [9], and E141 [10]. The ε values, which would explain the discrepancy between the measurement and the SM calculation of the anomalous magnetic moment of the muon [3] are displayed in Fig. 3 as the bold solid line with a 2σ band.

In conclusion, we perform a search for a dark photon in the mass range between 1.5 and $3.4 \text{ GeV}/c^2$, where we do not observe




Fig. 3. Exclusion limit at the 90% confidence level on the mixing parameter ε as a function of the dark photon mass. The bold solid line represents the ε values, which would explain the discrepancy between the measurement and the SM calculation of the anomalous magnetic moment of the muon [3], together with its 2σ band.

a significant signal. We set upper limits on the mixing parameter ε between 10^{-3} and 10^{-4} as a function of the dark photon mass with a confidence level of 90%. This is a competitive limit in this dark photon mass range. The BESIII results, which are based on two years of data taking, are already competitive to the large BaBar data samples, based on 9 years of running. This is possible due to the use of untagged ISR events for the dark photon search as well as the fact that the center-of-mass energy of the BEPCII collider is closer to the mass region tested. We also use a different analysis approach, which has no dependence on the radiator function.

The BESIII collaboration thanks the staff of BEPCII and the IHEP computing center for their strong support. This work is supported in part by National Key Basic Research Program of China under Contract No. 2015CB856700; National Natural Science Foundation of China (NSFC) under Contracts Nos. 11235011, 11335008, 11425524, 11625523, 11635010; the Chinese Academy of Sciences (CAS) Large-Scale Scientific Facility Program; the CAS Center for Excellence in Particle Physics (CCEPP); Joint Large-Scale Scientific Facility Funds of the NSFC and CAS under Contracts Nos. U1332201, U1532257, U1532258; CAS under Contracts Nos. KJCX2-YW-N29, KJCX2-YW-N45, QYZDJ-SSW-SLH003; 100 Talents Program of CAS; National 1000 Talents Program of China; INPAC and Shanghai Key Laboratory for Particle Physics and Cosmology; German Research Foundation DFG under Contracts Nos. Collaborative Research Center CRC 1044, FOR 2359; Istituto Nazionale di Fisica Nucleare, Italy; Joint Large-Scale Scientific Facility Funds of the NSFC and CAS; Koninklijke Nederlandse Akademie van Wetenschappen (KNAW) under Contract No. 530-4CDP03; Ministry of Development of Turkey under Contract No. DPT2006K-120470; National Natural Science Foundation of China (NSFC); National Science and Technology fund; The Swedish Research Council; U.S. Department of Energy under Contracts Nos. DE-FG02-05ER41374, DE-SC-0010118, DE-SC-0010504, DE-SC-0012069; University of Groningen (RuG) and the Helmholtzzentrum fuer Schwerionenforschung GmbH (GSI), Darmstadt; WCU Program of National Research Foundation of Korea under Contract No. R32-2008-000-10155-0.

References

- [1] N. Arkani-Hamed, D.P. Finkbeiner, T.R. Slatyer, N. Weiner, *Phys. Rev. D* 79 (2009) 015014.
- [2] B. Holdom, *Phys. Lett. B* 166 (1986) 196–198.
- [3] M. Pospelov, *Phys. Rev. D* 80 (2009) 095002.
- [4] J.D. Bjorken, R. Essig, P. Schuster, N. Toro, *Phys. Rev. D* 80 (2009) 075018.
- [5] B. Batell, M. Pospelov, A. Ritz, *Phys. Rev. D* 79 (2009) 115008.
- [6] H.B. Li, T. Luo, *Phys. Lett. B* 686 (2010) 249–253.
- [7] H. Merkel, et al., A1 Collaboration, *Phys. Rev. Lett.* 106 (2011) 251802.
- [8] H. Merkel, et al., A1 Collaboration, *Phys. Rev. Lett.* 112 (2014) 221802.
- [9] A. Bross, et al., E774 Collaboration, *Phys. Rev. Lett.* 67 (1991) 2942.
- [10] E.M. Riordan, et al., E141 Collaboration, *Phys. Rev. Lett.* 59 (1987) 755.
- [11] B. Aubert, et al., BaBar Collaboration, *Phys. Rev. Lett.* 103 (2009) 081803.
- [12] J.P. Lees, et al., BaBar Collaboration, *Phys. Rev. Lett.* 113 (2014) 201801.
- [13] M. Ablikim, et al., BESIII Collaboration, *Phys. Lett. B* 753 (2016) 629–638.
- [14] M. Ablikim, et al., BESIII Collaboration, *Nucl. Instr. Meth. A* 614 (2010) 345–399.
- [15] G. Rodrigo, H. Czyż, J.H. Kuhn, M. Szopa, *Eur. Phys. J. C* 24 (2002) 71.
- [16] H. Czyż, J.H. Kuhn, A. Wapient, *Phys. Rev. D* 77 (2008) 114005.
- [17] G. Balossini, C.M.C. Calame, G. Montagna, O. Nicosini, F. Piccinini, *Nucl. Phys. B* 758 (2006) 227–253.
- [18] S. Jadach, B.F.L. Ward, Z. Was, *Comput. Phys. Commun.* 130 (2000) 260–325.
- [19] J. Allison, et al., GEANT4 Collaboration, *IEEE Trans. Nucl. Sci.* 53 (2006) 270–278.
- [20] S. Agostinelli, et al., GEANT4 Collaboration, *Nucl. Instr. Meth. A* 506 (2003) 250–303.
- [21] D.M. Asner, et al., *Int. J. Mod. Phys. A* 24 (S1) (2009) 794.
- [22] V.P. Druzhinin, S.I. Eidelman, S.I. Serednyakov, E.P. Solodov, *Rev. Mod. Phys.* 83 (2011) 1545.
- [23] W.A. Rolke, A.M. Lopez, J. Conrad, *Nucl. Instrum. Methods Phys. Res., Sect. A* 551 (2005) 493–503.
- [24] G. Balossini, C. Bignamini, C.M.C. Calame, G. Montagna, O. Nicosini, F. Piccinini, *Phys. Lett. B* 663 (2008) 209–213.
- [25] T. Beranek, H. Merkel, M. Vanderhaeghen, *Phys. Rev. D* 88 (2013) 015032.
- [26] C. Patrignani, et al., Particle Data Group, *Chin. Phys. C* 40 (2016) 100001.
- [27] F. Archilli, et al., KLOE-2 Collaboration, *Phys. Lett. B* 706 (2012) 251–255.
- [28] D. Babuski, et al., KLOE-2 Collaboration, *Phys. Lett. B* 736 (2014) 459–464.
- [29] A. Anastasi, et al., KLOE-2 Collaboration, *Phys. Lett. B* 750 (2015) 633–637.
- [30] A. Anastasi, et al., KLOE-2 Collaboration, *Phys. Lett. B* 757 (2016) 356–361.
- [31] P. Adlarson, et al., WASA-at-COSY Collaboration, *Phys. Lett. B* 726 (2013) 187–193.
- [32] G. Agakishiev, et al., HADES Collaboration, *Phys. Lett. B* 731 (2014) 265–271.
- [33] A. Adare, et al., PHENIX Collaboration, *Phys. Rev. C* 91 (2015) 031901.
- [34] J.R. Batley, et al., NA48/2 Collaboration, *Phys. Lett. B* 746 (2015) 178–185.
- [35] S. Abrahamyan, et al., APEX Collaboration, *Phys. Rev. Lett.* 107 (2011) 191804.