

School of Natural Sciences and Mathematics

***Environmental and Genetic Determinants of
Biofilm Formation in Paracoccus Denitrificans***

UT Dallas Author(s):

Santosh Kumar
Stephen Spiro

Rights:

CC BY 4.0 (Attribution)
©2017 The Authors

Citation:

Kumar, Santosh, and Stephen Spiro. 2017. "Environmental and genetic determinants of biofilm formation in paracoccus denitrificans." *mSphere* 2(5), doi:10.1128/mSphereDirect.00350-17

This document is being made freely available by the Eugene McDermott Library of the University of Texas at Dallas with permission of the copyright owner. All rights are reserved under United States copyright law unless specified otherwise.



Environmental and Genetic Determinants of Biofilm Formation in *Paracoccus denitrificans*

Santosh Kumar,  Stephen Spiro

Department of Biological Sciences, University of Texas at Dallas, Richardson, Texas, USA

ABSTRACT The genome of the denitrifying bacterium *Paracoccus denitrificans* predicts the expression of a small heme-containing nitric oxide (NO) binding protein, H-NOX. The genome organization and prior work in other bacteria suggest that H-NOX interacts with a diguanylate cyclase that cyclizes GTP to make cyclic di-GMP (cdGMP). Since cdGMP frequently regulates attached growth as a biofilm, we first established conditions for biofilm development by *P. denitrificans*. We found that adhesion to a polystyrene surface is strongly stimulated by the addition of 10 mM Ca²⁺ to rich media. The genome encodes at least 11 repeats-in-toxin family proteins that are predicted to be secreted by the type I secretion system (TISS). We deleted the genes encoding the TISS and found that the mutant is almost completely deficient for attached growth. Adjacent to the TISS genes there is a potential open reading frame encoding a 2,211-residue protein with 891 Asp-Ala repeats. This protein is also predicted to bind calcium and to be a TISS substrate, and a mutant specifically lacking this protein is deficient in biofilm formation. By analysis of mutants and promoter reporter fusions, we show that biofilm formation is stimulated by NO generated endogenously by the respiratory reduction of nitrite. A mutant lacking both predicted diguanylate cyclases encoded in the genome overproduces biofilm, implying that cdGMP is a negative regulator of attached growth. Our data are consistent with a model in which there are H-NOX-dependent and -independent pathways by which NO stimulates biofilm formation.

IMPORTANCE The bacterium *Paracoccus denitrificans* is a model for the process of denitrification, by which nitrate is reduced to dinitrogen during anaerobic growth. Denitrification is important for soil fertility and greenhouse gas emission and in waste and water treatment processes. The ability of bacteria to grow as a biofilm attached to a solid surface is important in many different contexts. In this paper, we report that attached growth of *P. denitrificans* is stimulated by nitric oxide, an intermediate in the denitrification pathway. We also show that calcium ions stimulate attached growth, and we identify a large calcium binding protein that is required for growth on a polystyrene surface. We identify components of a signaling pathway through which nitric oxide may regulate biofilm formation. Our results point to an intimate link between metabolic processes and the ability of *P. denitrificans* to grow attached to a surface.

KEYWORDS biofilm, cyclic di-GMP, denitrification, flavohemoglobin, H-NOX, nitric oxide, *Paracoccus denitrificans*

Paracoccus denitrificans is a metabolically versatile alphaproteobacterium that has been extensively used as a model for studies of oxidative phosphorylation, anaerobic respiration (denitrification), and the metabolism of one-carbon compounds (1). Typically described as soil bacteria, *Paracoccus* spp. in fact seem rather ubiquitous, being also found in aquatic environments (2), waste treatment processes (3), and the

Received 7 August 2017 Accepted 16 August 2017 Published 6 September 2017

Citation Kumar S, Spiro S. 2017. Environmental and genetic determinants of biofilm formation in *Paracoccus denitrificans*. *mSphere* 2:e00350-17. <https://doi.org/10.1128/mSphereDirect.00350-17>.

Editor Katherine McMahon, University of Wisconsin—Madison

Copyright © 2017 Kumar and Spiro. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Stephen Spiro, stephen.spiro@utdallas.edu.

Solicited external reviewers: Elizabeth Boon, Stony Brook University; Andrew Gates, University of East Anglia.

This paper was submitted via the [mSphereDirect™](https://mspheredirect.org/) pathway.


 Santosh Kumar and @stephenspiro from @UTD_Biology explore biofilm formation and nitric oxide metabolism in *Paracoccus denitrificans*

TABLE 1 *Paracoccus denitrificans* genes discussed in this paper

Gene no.	Gene name	Function
Pden_0876	<i>pdeA</i>	cdGMP phosphodiesterase (predicted to be inactive)
Pden_1689	<i>hmp</i> (<i>fhp</i>)	NO denitrosylase
Pden_1690	<i>nsrR1</i>	NO-sensitive repressor NsrR
Pden_2025	<i>pdeB</i>	cdGMP phosphodiesterase
Pden_2411	<i>bapD</i>	TISS outer membrane protein
Pden_2412	<i>bapC</i>	TISS membrane fusion protein
Pden_2413	<i>bapB</i>	TISS ABC transporter
n/a ^a	<i>bapA</i>	TISS-secreted RTX family agglutinin
Pden_3024	<i>nsrR2</i>	NO-sensitive repressor NsrR
Pden_3025	<i>ytfE</i>	RIC family [Fe-S] repair protein
Pden_3719	<i>hnoX</i>	H-NOX protein
Pden_3720	<i>dgcA</i>	Diguanylate cyclase
Pden_3982	<i>dgcB</i>	Diguanylate cyclase (fused to response regulator domain)

^an/a, the *bapA* gene is not present in the original annotation of the *P. denitrificans* genome and so does not have a Pden gene number. The reading frame is present in the manually curated SEED annotation (<http://rast.nmpdr.org/seedviewer.cgi?page=Annotation&feature=figj318586.4.peg.2260>) and has been described by other authors (41).

deep subsurface (4) and in association with plants (5) and animals (6). *P. denitrificans* is a facultative anaerobe and an obligatorily respiratory organism. Under anoxic growth conditions, substrate oxidation is coupled to the sequential reduction of nitrogen oxyanions and oxides to dinitrogen, in the pathway called denitrification. Denitrification is an important route for the loss of fixed nitrogen from soil and is applied in water and waste treatment processes. In both cases, denitrification attracts interest as a significant source of the greenhouse gas nitrous oxide (7).

Nitric oxide (NO) metabolism by *P. denitrificans* is thought to depend principally upon a respiratory membrane-associated NO reductase, NOR (or NorBC), which couples oxidation of cytochrome *c* to the reduction of NO to nitrous oxide (N₂O). Expression of the *nor* operon is activated in response to NO by the transcriptional regulator NNR, which also activates genes encoding the respiratory nitrite reductase NirS that reduces nitrite to NO (8, 9). Thus, NO is a key regulatory signal controlling the transition from aerobic to anaerobic metabolism (10, 11).

The *P. denitrificans* genome reveals a rich repertoire of unstudied genes and proteins that potentially mediate additional responses to NO (Table 1). There are two homologs of the NO-sensitive repressor NsrR (12, 13). On the basis of genome context, one (Pden_1690) is predicted to regulate expression of the *hmp* gene (Pden_1689) encoding an NO-scavenging flavohemoglobin (14, 15), together with a closely linked gene of unknown function (Pden_1688). The second NsrR (Pden_3024) is predicted to regulate expression of *ytfE*, which encodes a protein from the RIC family that is involved in the repair of NO-damaged [Fe-S] proteins (16). Additionally, the genome encodes H-NOX (Pden_3719), a small NO binding heme protein (17–19). Biological roles of H-NOX proteins have not been well studied, but there are examples involved in the regulation of iron metabolism (20) and of motility versus attached growth (18, 21). The *P. denitrificans* H-NOX is encoded by a gene that overlaps the gene (Pden_3720) encoding a protein with a GGDEF domain. The GGDEF domain is associated with a diguanylate cyclase activity that cyclizes GTP to form cyclic di-GMP (cdGMP), a second messenger involved in the regulation of many bacterial processes (22). The *P. denitrificans* genome organization suggests that NO has the potential to regulate cdGMP synthesis and downstream processes, as is the case in some other bacteria (18, 19). In *Shewanella woodyi*, H-NOX interacts with a bifunctional diguanylate cyclase/cdGMP phosphodiesterase (23). In this case, the NO-bound form of H-NOX stimulates phosphodiesterase activity to lower the cytosolic concentration of cdGMP, an event which downregulates biofilm formation (24).

Biofilm formation and motility have been described as being reciprocally regulated by cdGMP in several species. Typically, cdGMP inhibits motility and stimulates attached growth as a biofilm (22). *P. denitrificans* is a nonmotile bacterium and lacks most




FIG 1 Assays of biofilm formation by *P. denitrificans* Pd1222 and by $\Delta bapBCD$ and $\Delta bapA$ mutants. Cultures were incubated for 72 h without shaking in 6-cm-diameter petri dishes. The growth medium was liquid L broth, supplemented with 10 mM calcium chloride as indicated. (A and B) Dishes were washed, stained with crystal violet, and photographed from the top (A) and side (B). (C) Crystal violet was extracted from biofilms and quantified by absorption at 595 nm. Means and standard deviations were calculated from results of triplicate assays. wt, wild type.

(though not all) genes that are associated with flagellum-based motility and chemotaxis. The genetic basis of biofilm formation by this organism has not been well studied but is a potential target for regulation by cdGMP. In biological water and sewage treatment processes, microorganisms frequently grow as surface-attached biofilms or as granules, and *Paracoccus* spp. have been observed in both cases (25, 26), suggesting that it is capable of attached growth. It is important to understand how metabolism in biofilms contributes to the efficiency of chemical transformations in treatment processes. In other bacteria, respiratory metabolism and electron acceptor availability are intimately linked to biofilm development (27).

In this paper, we report conditions that allow robust and reproducible biofilm formation by *P. denitrificans*. We show that attached growth is stimulated by calcium ions and suggest that attachment requires one or more calcium binding proteins secreted to the cell envelope through a type I secretion system. We present evidence to suggest that endogenously generated NO is a signal that stimulates attached growth and that the H-NOX/cdGMP signaling pathway has a role in the regulation of NO metabolism and biofilm formation.

RESULTS

Biofilm formation by *P. denitrificans*. We established conditions for biofilm formation in *P. denitrificans* and then assayed the phenotypes of mutants and the influence of environmental conditions. We found that *P. denitrificans* can form adhered biofilms when standing cultures are grown in rich medium in conventional polystyrene petri dishes. After 3 to 4 days of incubation, attached growth is detectable on the bottom surface of the petri dish (Fig. 1A), in addition to being detectable as a prominent ring that forms at the air-liquid interface (Fig. 1B). Bottom-attached growth is dependent on the surface area of the medium, being denser in 9-cm-diameter petri dishes than in 6-cm-diameter petri dishes and absent in 1-cm-diameter 24-well microplates (not shown). The effect of surface area and the growth at the air-liquid interface suggest that oxygen availability is a determinant of biofilm formation in *P. denitrificans*. Biofilm formation at the air-liquid interface occurs commonly, and this behavior has been observed in *Pseudomonas fluorescens* and *Escherichia coli* (28, 29), among other bacterial species.

Identification of a secretion pathway and potential adhesin required for attached growth. Sessile attached cells typically grow in a matrix that is formed by extracellular polysaccharide, nucleic acid, or protein. Information in the KEGG database (30) provides no evidence of an ability of *P. denitrificans* to synthesize extracellular polysaccharide. The genome annotation suggests that glycogen is the only polysaccharide made by this organism. We therefore searched for evidence of extracellular




FIG 2 Detection of BapA in envelope fractions of *P. denitrificans*. Planktonic phase cells were harvested from biofilm assays (grown in liquid L broth supplemented with 10 mM calcium chloride), and envelope fractions were prepared and separated by SDS-PAGE. The protein indicated by the arrow was shown by mass spectrometry to be BapA. The strains analyzed were as follows: lane 1, $\Delta bapBCD$ mutant; lane 2, wild type; lane 3, $\Delta hnoX$ mutant; lane 4, $\Delta dgcA$ mutant; lane 5, $\Delta dgcB$ mutant; lane 6, $\Delta dgcA \Delta dgcB$ mutant; lane 7, Δhmp mutant; lane 8, $\Delta hmp \Delta hnoX$ mutant; lane 9, Δnor mutant. Two gel images are shown since the number of samples exceeded the capacity of the gel apparatus.

proteins that might function as adhesins. We used PSORTdb (31) to retrieve predictions of the subcellular location of the *P. denitrificans* proteome. This analysis identified 11 extracellular or cell envelope-associated proteins from the repeats-in-toxin (RTX) family (see Fig. S1 in the supplemental material). These proteins of diverse function are characterized by the presence of multiple nonapeptide calcium binding domains and secretion by the type I secretion system (32). The RTX family includes proteins that function as adhesins (33). In 3-day incubations, we found that the addition of 10 mM Ca^{2+} to growth media strongly stimulated biofilm formation (Fig. 1). These observations are consistent with the idea of attached growth depending on extracellular Ca^{2+} binding proteins. The calcium ion concentration of a rich medium similar to ours has been estimated to be 120 μM (34). Our defined medium formulation contains 0.68 mM calcium chloride and 19.6 μM EDTA (35). Compared to the results seen with rich medium, we observed more attached growth in unsupplemented minimal medium (consistent with the higher calcium concentration) and so a lower level of stimulation by additional calcium (data not shown). Calcium supplementation was also prone to causing precipitation in minimal medium. Therefore, all cultures discussed here were grown in rich medium in 6-cm-diameter petri dishes.

We analyzed the *P. denitrificans* proteome using the TXSScan algorithm (36), which revealed the presence of a single predicted type I secretion system (TISS) encoded by Pden_2411 to Pden_2413 (Table 1). These genes encode the ABC transporter (Pden_2413), membrane fusion protein (Pden_2412), and outer membrane efflux protein (Pden_2411) components of the TISS (37). We designate these genes *bapBCD*, for reasons explained below. To generate a null TISS-deficient mutant, we deleted these three genes. The resultant $\Delta bapBCD$ strain had no gross growth defect in liquid culture or on agar (not shown) but was almost completely deficient for biofilm formation on polystyrene (Fig. 1). Our minimal interpretation of these results is that attached growth in *P. denitrificans* requires one or more of the extracellular Ca^{2+} binding proteins that are secreted by the TISS pathway.

To search for direct evidence of extracellular adhesins, cells from the planktonic phase of biofilm assays (grown in rich medium supplemented with 10 mM Ca^{2+}) were fractionated to enrich for periplasmic and outer membrane proteins (38), which were then separated by denaturing polyacrylamide gel electrophoresis (SDS-PAGE). A very large (>300-kDa) protein was visible in envelope fractions (Fig. 2) and had increased abundance in mutants with hyperbiofilm phenotypes (see below). The accumulation of this protein correlates very well with genotypes that stimulate biofilm accumulation, which, together with its size and location, makes it a good candidate for an adhesin.

In the manually curated SEED (39) annotation of the *P. denitrificans* genome, there is a 6,636-bp gene immediately upstream of the three genes encoding the TISS (<http://rast.nmpdr.org/seedviewer.cgi?page=Annotation&feature=fig|318586.4.peg.2260>). The 2,211-amino-acid product of this gene (designated *bapA*, as explained below) is annotated as a TISS-secreted RTX family agglutinin. A total of 1,782 residues of the BapA protein consist of 891 repeats of the dipeptide Asp-Ala, with divergent

sequences at the N and C termini. The PSORTdb algorithm (31) strongly predicts BapA to be located in the outer membrane (it was absent from our original analysis because this open reading frame is annotated only in the SEED database). The BapA protein has a molecular mass of 212 kDa and a predicted pI of 2.4. Mass spectrometry (MS) of excised gel slices confirmed that the largest envelope protein detected by SDS-PAGE is BapA. In two independent samples, BapA was identified as the most abundant protein in the excised band by normalized spectral counts and the spectral index criterion (40). All BapA tryptic peptides below 40 amino acids in length were detected, and BapA was identified with 134 peptide spectrum matches and 10 unique peptide sequences, with a false-discovery rate of 1%. Thus, we have confidence in the assignment of the stained band in envelope fractions (Fig. 2) as BapA. The BapA protein has an apparent molecular weight in SDS-PAGE that is significantly higher than its true molecular weight (Fig. 2). This discrepancy may relate to the highly unusual amino acid composition of BapA (41.8% alanine, 42.6% aspartate), although we cannot exclude the possibility of a posttranslational modification. We constructed an unmarked deletion in the *bapA* gene and found that the mutant has a phenotype similar to that of the strain lacking the TISS, that is, a severe defect in biofilm formation (Fig. 1). BapA is almost undetectable on Coomassie-stained gels of the envelope fraction of a wild-type strain (Fig. 2, lane 2). Nevertheless, we assume that the wild-type strain makes sufficient BapA to promote the degree of attached growth visible in petri dishes.

To summarize, we have shown that biofilm formation by *P. denitrificans* is stimulated by calcium ions and depends upon a functional TISS. Genetic evidence implicates a large envelope protein that is a predicted TISS substrate as being required for attached growth, perhaps functioning as an adhesin. We cannot exclude the possibility that other TISS substrates (of which there are at least 11 predicted) also have roles in attached growth.

As we were completing this work, similar conclusions about the role of the *P. denitrificans* TISS and the RTX family agglutinin were reported elsewhere (41). The three genes encoding the components of the TISS were designated *bapBCD*, and the gene encoding the secreted adhesion was designated *bapA*; we have therefore adopted this nomenclature (Table 1). Our work extends the findings of Yoshida et al. (41) by demonstrating a clear role for calcium ions in attached growth, by detecting BapA directly by SDS-PAGE, and by showing that BapA abundance correlates, at least qualitatively, with the extent of attached growth (see below).

Endogenously generated nitric oxide stimulates biofilm formation. Recent work has shown that the flavohemoglobin of *P. denitrificans* protects against extracellular nitrite during aerobic growth, presumably because of its ability to scavenge traces of NO generated abiotically from nitrous acid (42). We have constructed an *hmp* mutant and confirmed that it metabolizes NO more slowly than the wild type in whole-cell assays of NO consumption (data not shown). Surprisingly, the *hmp* mutant also overproduces biofilm compared to a wild-type strain (Fig. 3). We have previously shown that our formulation of L broth contains approximately 0.5 mM nitrate which can act as a source of endogenously generated NO under oxygen-limiting conditions (43). To explain the phenotype of the *hmp* mutant, we hypothesize that oxygen limitation in these static cultures initiates expression of denitrification genes. Thus, nitrate is reduced to nitrite and nitrite to NO. Since the *hmp* mutant is lacking one pathway for NO consumption, elevated NO concentrations in this strain might explain the increase in biofilm, if NO is a signal that stimulates biofilm formation. To test this hypothesis, we first grew static biofilm cultures of strains containing *hmp-lacZ* and *norC-lacZ* reporter fusions. The *norC* promoter is activated by the NO-sensing transcription factor NNR (44), and the *hmp* promoter is predicted to be subject to negative regulation by the NO-sensitive repressor NsrR1. In static cultures in unsupplemented medium, the *hmp* promoter has activity as high as that measured in cultures grown in shake flasks and treated with a source of NO (Table 2). In contrast, the *norC* promoter has a rather low and submaximal level of activity in unsupplemented static cultures. Therefore, we




FIG 3 Assays of biofilm formation by *P. denitrificans* Pd1222 and $\Delta hnoX$, $\Delta dgcA$, $\Delta dgcB$, $\Delta dgcAB$, Δhmp , and Δnor mutants. Cultures were incubated for 72 h without shaking in 6-cm-diameter petri dishes. The growth medium was L broth, supplemented with 10 mM calcium chloride and 5 mM nitrate as indicated. (A) Dishes were washed, stained with crystal violet, and photographed from the top and side. (B) The crystal violet was extracted with ethanol, and the absorbance was measured at 595 nm (68). Means and standard deviations were calculated from results of triplicate assays. The strains used were either wild type (*nirS*⁺) or deleted ($\Delta nirS$) for the nitrite reductase gene *nirS* as indicated.

conclude that NO is generated endogenously in static cultures to an extent that efficiently induces the *hmp* but not the *norC* promoter.

To confirm the source of the endogenously generated NO and that it stimulates biofilm formation, we introduced a *nirS* mutation into the previously characterized strains. Mutation of *nirS* abolishes biofilm formation in both the wild-type strain and the *hmp* mutant (Fig. 3), which strongly suggests that reduction of nitrite to NO generates a signal that stimulates biofilm formation. It is unclear why the $\Delta nirS \Delta nor$ double mutant overproduces biofilm compared to the $\Delta nirS$ strain (Fig. 3).

The abundance of BapA is higher in the *hmp* mutant than in the wild-type strain (Fig. 2, lane 7), which may explain the hyperbiofilm phenotype of the *hmp* mutant. An interesting possibility is that elevated NO levels in the *hmp* mutant might lead (either directly or indirectly) to increased expression of the *bapA* gene.

Regulation of biofilm formation by cdGMP. According to the MIST database (45), the *P. denitrificans* genome encodes two proteins with GGDEF domains associated with

TABLE 2 Activities of promoter reporter fusions measured in shake flasks and in static cultures

Reporter fusion	β -Galactosidase activity (mean \pm SD) ^a			
	Shake flask	Shake flask + NO	Static	Static + 5 mM nitrate
<i>hmp-lacZ</i>	391 \pm 12	824 \pm 12	895 \pm 40	1,205 \pm 109
<i>norC-lacZ</i>	91 \pm 5	329 \pm 17	179 \pm 5	2,103 \pm 140

^aLiquid cultures were grown in shake flasks to mid-log phase or for 72 h in static 6-cm-diameter petri dishes (the same conditions used for biofilm assays). For NO-treatment, shake flask cultures were amended with 50 μ M spermine NONOate 60 min prior to sampling. β -Galactosidase was assayed according to Miller (69). Means and standard deviations were calculated from results of triplicate assays.




FIG 4 A model for the regulation of biofilm formation in *Paracoccus denitrificans* based on the work described in this paper. Abbreviations: H-NOX, heme NO and oxygen binding; REC, receiver domain; HK, histidine kinase. GGDEF and EAL are sequence motifs associated with the diguanylate cyclase and phosphodiesterase, respectively. Enzymes and enzyme-catalyzed reactions are shown in black, signaling interactions in blue and gray (the latter indicating pathways that are likely to be indirect). We suggest that Hmp is the dominant pathway for NO metabolism in the static cultures used for biofilm assays.

the synthesis of cdGMP (Fig. 4). One (Pden_3720; DgcA) has an N-terminal domain of unknown function, and the product of the adjacent Pden_3719 gene is an H-NOX protein (18). Like the H-NOX from its close relative *Rhodobacter sphaeroides* (17), the *P. denitrificans* H-NOX lacks the otherwise conserved histidine residue which functions as the axial ligand to the heme (18). The genome organization suggests that H-NOX regulates the biosynthesis of cdGMP. There is a similar situation in *Legionella pneumophila* and *Shewanella woodyi*, though in these cases the H-NOX protein regulates the activities of a bifunctional enzyme that both synthesizes and degrades cdGMP (24, 46). The second GGDEF domain-containing protein (Pden_3982; DgcB) has an N-terminal receiver domain, implying a signal input pathway from an unidentified histidine kinase. The *P. denitrificans* genome encodes two proteins with EAL domains associated with cdGMP hydrolysis (Fig. 4), but one of these is predicted to be catalytically inactive (47). Catalytically inactive EAL domains can potentially function as cdGMP receptors (22).

To begin to explore the function of the cdGMP signaling pathway in *P. denitrificans*, we have characterized mutants lacking single and multiple genes encoding pathway components. In assays of nitric oxide (NO) consumption by washed cell suspensions, we found that the cdGMP-deficient mutant ($\Delta dgcAB$) consumes NO somewhat more slowly than the wild-type parent and that the *hnoX* mutant consumes NO more rapidly (Fig. 5). Reporter fusion assays showed no effects of *hnoX* and *dgcAB* mutations on *norC* or *hmp* promoter activity (data not shown). Therefore, it is likely that cdGMP regulation of NO consumption does not operate at the level of gene expression.

In biofilm assays, the *hnoX* mutant is partially deficient for attached growth, while the *dgcAB* and *hmp* mutants overproduce biofilm (Fig. 3). The BapA protein overaccumulates in the same strains (mutants Δhmp and $\Delta dgcAB$) that overproduce biofilm and is absent in the *hnoX* mutant (Fig. 2, lanes 7, 6, and 3, respectively), further evidence of a correlation between biofilm density and BapA abundance, which is consistent with this protein having a function in attached growth. The data also suggest that *bapA* expression or the secretion of the BapA protein to the cell envelope (or both) is regulated by cdGMP and NO.

The overproduction of biofilm by the *dgcAB* mutant is consistent with the possibility that cdGMP regulates biofilm formation and/or that NO accumulates in this mutant (because of the reduced rate of consumption) and stimulates biofilm formation as described above. To distinguish these possibilities, we introduced the $\Delta nirS$ mutation into the *dgcAB* mutant. The *dgcAB nirS* mutant continued to overproduce biofilm




FIG 5 Assays of NO consumption by washed cell suspensions of *P. denitrificans* and its *hnoX* and *dgcAB* mutants. Cultures were grown aerobically in succinate minimal media (A) and were treated with 50 μ M spermine NONOate for 60 min prior to harvesting (B) in order to induce expression of the *hmp* and *nor* genes.

compared to the $\Delta nirS$ parent (Fig. 3). This observation implies that the increased-biofilm phenotype of the *dgcAB* mutant cannot be ascribed solely to accumulation of NO and that cdGMP may therefore negatively regulate biofilm formation independently of NO (Fig. 4).

DISCUSSION

Large cell surface proteins that have roles in the biofilm matrix have been identified in multiple species (33, 48–50). In many cases, these proteins are characterized by the occurrence of complex and/or simple repeat sequences. One such protein that is quite well understood and that may serve as a model for BapA is SdrC of *Staphylococcus aureus* (51). SdrC is a 995-residue multidomain protein with 109 Ser-Asp repeats. SdrC is involved in cell-cell and cell-surface interactions, and these appear to be separable functions of the protein. Homophilic SdrC-mediated cell-cell interactions require two pentapeptide sequences, RPGSV and VDQYT, separated by 37 residues (51). Interestingly, there are very similar sequences (LPGSV and VDRYT) in the *P. denitrificans* BapA protein, although the spacing is much greater, since these motifs occur on either side of the Asp-Ala repeats. SdrC also promotes adhesion to plastic surfaces (51); this is a strong interaction believed to involve the unfolding of SdrC, exposing hydrophobic residues responsible for the protein-surface adhesion (52). The N- and C-terminal domains of BapA that flank the Asp-Ala repeats are quite hydrophobic (being 26% and 22% aliphatic residues, respectively) and have sequence similarity to other adhesins (41). Thus, it is possible that BapA mediates both cell-cell and cell-surface interactions in a manner similar to that seen with SdrC. This idea is consistent with the observation that BapA contributes to cell surface hydrophobicity (41), although the suggestion that the Asp-Ala repeats contribute directly to hydrophobicity (41) does not seem intuitively reasonable. The function of the Ser-Asp repeats of SdrC is not known; nor is the role of metal ions in SdrC function fully understood. SdrC-mediated biofilm formation is inhibited by manganese ions but is not affected by calcium (51).

The primary structure of BapA and the stimulatory effect of calcium ions suggest other possible mechanisms for BapA activity. Extracellular aspartate-rich proteins that bind calcium ions have been documented in mammals (53, 54). Aspartate-rich proteins are also found in the shell matrices of bivalves such as oysters. Aspein is one such protein and is characterized by extreme aspartate richness and the presence of Asp-Ala repeat sequences (55). Cell surface carboxylates may function to nucleate the precipitation of calcium carbonate (56), and aspein and other polyanionic proteins are believed to play a role in the formation of the calcite mineral component of the shell (57). Bacteria are well known to mineralize calcium, and denitrification is one of the

metabolic processes associated with calcium carbonate formation (58). Calcium carbonate has been detected in bacterial biofilms, for example, in the case of *Pseudomonas aeruginosa* (59); thus, we speculate that BapA might function to nucleate the formation of a calcium mineral that is required for cell-cell and cell-surface interactions. An alternative possibility is that binding of calcium ions to BapA alters the structure of the protein in a way that is required for it to function as an adhesin. This mechanism operates for some other large cell surface calcium-dependent adhesins, such as LapF of *Pseudomonas fluorescens* (60).

Our mutant analysis suggests a role for the H-NOX pathway and cdGMP in the regulation of biofilm formation (Fig. 4). In contrast to most other bacteria (22), cdGMP acts negatively on biofilm, and this can, at least in part, be attributed to regulation of BapA abundance (Fig. 1 and 2). We suggest that H-NOX inhibits the activity of DgcA, since the *hnoX* mutant underproduces both biofilm and BapA (Fig. 2 and 3). This implies that NO stimulates H-NOX-mediated inhibition of DgcA to reduce the size of the pool of cdGMP and so to stimulate biofilm formation. Given that NirS-derived NO stimulates biofilm in *hnoX* and *dgc* mutants (Fig. 3) and that an *hnoX hmp* double mutant forms biofilm and overproduces BapA like an *hmp* mutant (Fig. 2 and data not shown), we believe that NO can also stimulate biofilm formation independently of H-NOX/DgcA (Fig. 4).

MATERIALS AND METHODS

Bacterial strains and growth conditions. All *P. denitrificans* mutants were derived from strain Pd1222 (61). *Escherichia coli* S17-1 (62) was used as the donor for the introduction of plasmids into *P. denitrificans* by conjugation (63). *E. coli* Novablue Singles competent cells (Novagen) were used as the host for routine DNA manipulations. The rich medium was L broth (10 g tryptone-liter⁻¹, 5 g yeast extract-liter⁻¹, 10 g NaCl-liter⁻¹), and the defined medium for *P. denitrificans* was as previously described (35). Bacteria were cultured at 37°C (*E. coli*) or 30°C (*P. denitrificans*).

Genetic manipulations. Unmarked deletions were made by allelic replacement using pK18*mobsacB* (64) and the method described by Sullivan et al. (65). Briefly, in-frame unmarked deletions were constructed by PCR amplification of ~600-bp fragments from each side of the gene of interest. The two PCR amplicons were ligated together by PCR using overlapping regions added in the primers (66). The ligated fragment was cloned into pK18*mobsacB*, transformed into *E. coli* S17-1 and then mobilized into *P. denitrificans* by conjugation. Single-crossover events were selected on kanamycin plates, and then recombinants with double crossovers were isolated on the basis of sucrose resistance and kanamycin sensitivity. The structures of chromosomal deletions were confirmed by PCR. Promoter-*lacZ* reporter fusions were constructed by cloning ~400 bp upstream of the start codon of genes of interest into pMP220 (67). Plasmid clones were introduced into *P. denitrificans* by conjugation from *E. coli* S17-1. Triparental matings were performed by mixing log-phase cells of *P. denitrificans*, *E. coli* S17-1 cells harboring recombinant pMP220 or pK18*mobsacB* derivatives, and *E. coli* DH5 α (pRK2013) cells in a 2:1:1 ratio and by incubation on L agar for 24 h at 30°C. *P. denitrificans* exconjugants were selected by analysis of rifampin or tetracycline resistance. Plasmid pSTBlue-1 (Novagen) was used as the vector for routine DNA manipulations and sequencing.

Biofilm assays. Attached growth was assayed in 60-mm-by-15-mm polystyrene petri dishes. A 100- μ l aliquot of overnight culture was inoculated into 7 ml of L broth amended with calcium chloride as indicated. The petri dishes were incubated for about 72 h at 30°C without shaking. The plates were then washed three times with water and treated with crystal violet (0.1% in water) for 20 min, washed three times with water, and air dried. For quantitation, the stain was extracted into 2 ml 95% ethanol and diluted as necessary, and the absorbance was measured at 595 nm (68).

Biochemical and other methods. Envelope proteins were enriched by cell fractionation (38) and resolved by denaturing gel electrophoresis using 5% polyacrylamide. Stained bands of interest were excised from gels and analyzed by mass spectrometry in the Proteomics Facility at the University of Texas Southwestern Medical School. Tandem mass spectrometry (MS/MS) spectra were collected for tryptic fragments, and data were analyzed with an in-house pipeline, with quantitation according to the normalized spectral index method (40). Beta-galactosidase activity was measured in cells made permeable by treatment with chloroform and SDS (69). Nitric oxide consumption by washed cell suspensions (normalized according to optical density) was measured with an amperometric NO-specific electrode (ISO NOP electrode; WPI Instruments) as described previously (70).

SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <https://doi.org/10.1128/mSphereDirect.00350-17>.

FIG S1, EPS file, 2.2 MB.

ACKNOWLEDGMENTS

We are grateful to Andrew Lemoff (University of Texas Southwestern Medical School) for his help with the collection and interpretation of mass spectrometry data.

This work was supported in part by the C. L. and Amelia A. Lundell endowment.

REFERENCES

- Baker SC, Ferguson SJ, Ludwig B, Page MD, Richter OM, van Spanning RJ. 1998. Molecular genetics of the genus *Paracoccus*: metabolically versatile bacteria with bioenergetic flexibility. *Microbiol Mol Biol Rev* 62: 1046–1078.
- Du H, Jiao N, Hu Y, Zeng Y. 2006. Diversity and distribution of pigmented heterotrophic bacteria in marine environments. *FEMS Microbiol Ecol* 57:92–105. <https://doi.org/10.1111/j.1574-6941.2006.00090.x>.
- Wang Z, Zhang XX, Lu X, Liu B, Li Y, Long C, Li A. 2014. Abundance and diversity of bacterial nitrifiers and denitrifiers and their functional genes in tannery wastewater treatment plants revealed by high-throughput sequencing. *PLoS One* 9:e113603. <https://doi.org/10.1371/journal.pone.0113603>.
- Russell JA, León-Zayas R, Wrighton K, Biddle JF. 2016. Deep subsurface life from North Pond: enrichment, isolation, characterization and genomes of heterotrophic bacteria. *Front Microbiol* 7:678. <https://doi.org/10.3389/fmicb.2016.00678>.
- Dias ACF, Dini-Andreote F, Hannula SE, Andreote FD, Pereira E Silva MDC, Salles JF, de Boer W, van Veen J, van Elsas JD. 2013. Different selective effects on rhizosphere bacteria exerted by genetically modified versus conventional potato lines. *PLoS One* 8:e67948. <https://doi.org/10.1371/journal.pone.0067948>.
- Castelino M, Eyre S, Moat J, Fox G, Martin P, Ho P, Upton M, Barton A. 2017. Optimisation of methods for bacterial skin microbiome investigation: primer selection and comparison of the 454 versus MiSeq platform. *BMC Microbiol* 17:23. <https://doi.org/10.1186/s12866-017-0927-4>.
- Richardson D, Felgate H, Watmough N, Thomson A, Baggs E. 2009. Mitigating release of the potent greenhouse gas N₂O from the nitrogen cycle—could enzymic regulation hold the key? *Trends Biotechnol* 27: 388–397. <https://doi.org/10.1016/j.tibtech.2009.03.009>.
- Van Spanning RJ, De Boer AP, Reijnders WN, Spiro S, Westerhoff HV, Stouthamer AH, Van der Oost J. 1995. Nitrite and nitric oxide reduction in *Paracoccus denitrificans* is under the control of NNR, a regulatory protein that belongs to the FNR family of transcriptional activators. *FEBS Lett* 360:151–154. [https://doi.org/10.1016/0014-5793\(95\)00091-M](https://doi.org/10.1016/0014-5793(95)00091-M).
- Van Spanning RJ, De Boer AP, Reijnders WN, Westerhoff HV, Stouthamer AH, Van Der Oost J. 1997. FnrP and NNR of *Paracoccus denitrificans* are both members of the FNR family of transcriptional activators but have distinct roles in respiratory adaptation in response to oxygen limitation. *Mol Microbiol* 23:893–907. <https://doi.org/10.1046/j.1365-2958.1997.2801638.x>.
- Bergaust L, Mao Y, Bakken LR, Frostegård A. 2010. Denitrification response patterns during the transition to anoxic respiration and post-transcriptional effects of suboptimal pH on nitrous oxide reductase in *Paracoccus denitrificans*. *Appl Environ Microbiol* 76:6387–6396. <https://doi.org/10.1128/AEM.00608-10>.
- Hassan J, Bergaust LL, Wheat ID, Bakken LR. 2014. Low probability of initiating *nirS* transcription explains observed gas kinetics and growth of bacteria switching from aerobic respiration to denitrification. *PLoS Comput Biol* 10:e1003933. <https://doi.org/10.1371/journal.pcbi.1003933>.
- Bodenmiller DM, Spiro S. 2006. The *yjeB* (*nsrR*) gene of *Escherichia coli* encodes a nitric oxide-sensitive transcriptional regulator. *J Bacteriol* 188:874–881. <https://doi.org/10.1128/JB.188.3.874-881.2006>.
- Tucker NP, Le Brun NE, Dixon R, Hutchings MI. 2010. There's NO stopping NsrR, a global regulator of the bacterial NO stress response. *Trends Microbiol* 18:149–156. <https://doi.org/10.1016/j.tim.2009.12.009>.
- Poole RK, Hughes MN. 2000. New functions for the ancient globin family: bacterial responses to nitric oxide and nitrosative stress. *Mol Microbiol* 36:775–783. <https://doi.org/10.1046/j.1365-2958.2000.01889.x>.
- Hausladen A, Gow A, Stamler JS. 2001. Flavohemoglobin denitrosylase catalyzes the reaction of a nitroxyl equivalent with molecular oxygen. *Proc Natl Acad Sci U S A* 98:10108–10112. <https://doi.org/10.1073/pnas.181199698>.
- Justino MC, Baptista JM, Saraiva LM. 2009. Di-iron proteins of the Ric family are involved in iron-sulfur cluster repair. *Biometals* 22:99–108. <https://doi.org/10.1007/s10534-008-9191-2>.
- Iyer LM, Anantharaman V, Aravind L. 2003. Ancient conserved domains shared by animal soluble guanylyl cyclases and bacterial signaling proteins. *BMC Genomics* 4:5. <https://doi.org/10.1186/1471-2164-4-5>.
- Plate L, Marletta MA. 2013. Nitric oxide-sensing H-NOX proteins govern bacterial communal behavior. *Trends Biochem Sci* 38:566–575. <https://doi.org/10.1016/j.tibs.2013.08.008>.
- Nisbett LM, Boon EM. 2016. Nitric oxide regulation of H-NOX signaling pathways in bacteria. *Biochemistry* 55:4873–4884. <https://doi.org/10.1021/acs.biochem.6b00754>.
- Wang Y, Dufour YS, Carlson HK, Donohue TJ, Marletta MA, Ruby EG. 2010. H-NOX-mediated nitric oxide sensing modulates symbiotic colonization by *Vibrio fischeri*. *Proc Natl Acad Sci U S A* 107:8375–8380. <https://doi.org/10.1073/pnas.1003571107>.
- Arora DP, Hossain S, Xu Y, Boon EM. 2015. Nitric oxide regulation of bacterial biofilms. *Biochemistry* 54:3717–3728. <https://doi.org/10.1021/bi501476n>.
- Römmling U, Galperin MY, Gomelsky M. 2013. Cyclic di-GMP: the first 25 years of a universal bacterial second messenger. *Microbiol Mol Biol Rev* 77:1–52. <https://doi.org/10.1128/MMBR.00043-12>.
- Lahiri T, Luan B, Raleigh DP, Boon EM. 2014. A structural basis for the regulation of an H-NOX-associated cyclic-di-GMP synthase/phosphodiesterase enzyme by nitric oxide-bound H-NOX. *Biochemistry* 53:2126–2135. <https://doi.org/10.1021/bi401597m>.
- Liu N, Xu Y, Hossain S, Huang N, Coursolle D, Gralnick JA, Boon EM. 2012. Nitric oxide regulation of cyclic di-GMP synthesis and hydrolysis in *Shewanella woodyi*. *Biochemistry* 51:2087–2099. <https://doi.org/10.1021/bi201753f>.
- Cydzik-Kwiatkowska A. 2015. Bacterial structure of aerobic granules is determined by aeration mode and nitrogen load in the reactor cycle. *Bioresour Technol* 181:312–320. <https://doi.org/10.1016/j.biortech.2015.01.101>.
- Singh S, Nerurkar AS, Srinandan CS. 2015. Nitrate levels modulate the abundance of *Paracoccus* sp. in a biofilm community. *World J Microbiol Biotechnol* 31:951–958. <https://doi.org/10.1007/s11274-015-1849-7>.
- Dietrich LEP, Okegbe C, Price-Whelan A, Sakhtah H, Hunter RC, Newman DK. 2013. Bacterial community morphogenesis is intimately linked to the intracellular redox state. *J Bacteriol* 195:1371–1380. <https://doi.org/10.1128/JB.02273-12>.
- O'Toole GA, Kolter R. 1998. Initiation of biofilm formation in *Pseudomonas fluorescens* WCS365 proceeds via multiple, convergent signalling pathways: a genetic analysis. *Mol Microbiol* 28:449–461. <https://doi.org/10.1046/j.1365-2958.1998.00797.x>.
- Colón-González M, Méndez-Ortiz MM, Membrillo-Hernández J. 2004. Anaerobic growth does not support biofilm formation in *Escherichia coli* K-12. *Res Microbiol* 155:514–521. <https://doi.org/10.1016/j.resmic.2004.03.004>.
- Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M. 2016. KEGG as a reference resource for gene and protein annotation. *Nucleic Acids Res* 44:D457–D462. <https://doi.org/10.1093/nar/gkv1070>.
- Peabody MA, Laird MR, Vlasschaert C, Lo R, Brinkman FSL. 2016. PSORTdb: expanding the bacteria and archaea protein subcellular localization database to better reflect diversity in cell envelope structures. *Nucleic Acids Res* 44:D663–D668. <https://doi.org/10.1093/nar/gkv1271>.
- Linhartová I, Bumba L, Mašín J, Basler M, Osička R, Kamanová J, Procházková K, Adkins I, Hejnová-Holubová J, Sadílková L, Morová J, Sebo P. 2010. RTX proteins: a highly diverse family secreted by a common mechanism. *FEMS Microbiol Rev* 34:1076–1112. <https://doi.org/10.1111/j.1574-6976.2010.00231.x>.
- Satchell KJF. 2011. Structure and function of MARTX toxins and other large repetitive RTX proteins. *Annu Rev Microbiol* 65:71–90. <https://doi.org/10.1146/annurev-micro-090110-102943>.
- Boehm DF, Welch RA, Snyder IS. 1990. Calcium is required for binding of *Escherichia coli* hemolysin (HlyA) to erythrocyte membranes. *Infect Immun* 58:1951–1958.
- Hahnke SM, Moosmann P, Erb TJ, Strous M. 2014. An improved medium

- for the anaerobic growth of *Paracoccus denitrificans* Pd1222. *Front Microbiol* 5:18. <https://doi.org/10.3389/fmicb.2014.00018>.
36. Abby SS, Cury J, Guglielmini J, Néron B, Touchon M, Rocha EPC. 2016. Identification of protein secretion systems in bacterial genomes. *Sci Rep* 6:23080. <https://doi.org/10.1038/srep23080>.
 37. Thomas S, Holland IB, Schmitt L. 2014. The type 1 secretion pathway—the hemolysin system and beyond. *Biochim Biophys Acta* 1843:1629–1641. <https://doi.org/10.1016/j.bbamcr.2013.09.017>.
 38. Quan S, Hiniker A, Collet JF, Bardwell JCA. 2013. Isolation of bacteria envelope proteins. *Methods Mol Biol* 966:359–366. https://doi.org/10.1007/978-1-62703-245-2_22.
 39. Overbeek R, Begley T, Butler RM, Choudhuri JV, Chuang HY, Cohoon M, de Crécy-Lagard V, Diaz N, Disz T, Edwards R, Fonstein M, Frank ED, Gerdes S, Glass EM, Goessmann A, Hanson A, Iwata-Reuyl D, Jensen R, Jamshidi N, Krause L, Kubal M, Larsen N, Linke B, McHardy AC, Meyer F, Newweger H, Olsen G, Olson R, Osterman A, Portnoy V, Pusch GD, Rodionov DA, Rückert C, Steiner J, Stevens R, Thiele I, Vassieva O, Ye Y, Zagnitko O, Vonstein V. 2005. The subsystems approach to genome annotation and its use in the project to annotate 1000 genomes. *Nucleic Acids Res* 33:5691–5702. <https://doi.org/10.1093/nar/gki866>.
 40. Trudgian DC, Ridlova G, Fischer R, Mackeen MM, Ternette N, Acuto O, Kessler BM, Thomas B. 2011. Comparative evaluation of label-free SINQ normalized spectral index quantitation in the central proteomics facilities pipeline. *Proteomics* 11:2790–2797. <https://doi.org/10.1002/pmic.201000800>.
 41. Yoshida K, Toyofuku M, Obana N, Nomura N. 2017. Biofilm formation by *Paracoccus denitrificans* requires a type I secretion system-dependent adhesin BapA. *FEMS Microbiol Lett* 364. <https://doi.org/10.1093/femsle/fnx029>.
 42. Hartop KR, Sullivan MJ, Giannopoulos G, Gates AJ, Bond PL, Yuan Z, Clarke TA, Rowley G, Richardson DJ. 2017. The metabolic impact of extracellular nitrite on aerobic metabolism of *Paracoccus denitrificans*. *Water Res* 113:207–214. <https://doi.org/10.1016/j.watres.2017.02.011>.
 43. Hutchings MI, Shearer N, Wastell S, van Spanning RJ, Spiro S. 2000. Heterologous NNR-mediated nitric oxide signaling in *Escherichia coli*. *J Bacteriol* 182:6434–6439. <https://doi.org/10.1128/JB.182.22.6434-6439.2000>.
 44. Hutchings MI, Spiro S. 2000. The nitric oxide regulated *nor* promoter of *Paracoccus denitrificans*. *Microbiology* 146:2635–2641. <https://doi.org/10.1099/00221287-146-10-2635>.
 45. Ulrich LE, Zhulin IB. 2007. MiST: a microbial signal transduction database. *Nucleic Acids Res* 35:D386–D390. <https://doi.org/10.1093/nar/gkl932>.
 46. Carlson HK, Vance RE, Marletta MA. 2010. H-NOX regulation of c-di-GMP metabolism and biofilm formation in *Legionella pneumophila*. *Mol Microbiol* 77:930–942. <https://doi.org/10.1111/j.1365-2958.2010.07259.x>.
 47. Seshasayee ASN, Fraser GM, Luscombe NM. 2010. Comparative genomics of cyclic-di-GMP signalling in bacteria: post-translational regulation and catalytic activity. *Nucleic Acids Res* 38:5970–5981. <https://doi.org/10.1093/nar/gkq382>.
 48. Speziale P, Pietrocola G, Foster TJ, Geoghegan JA. 2014. Protein-based biofilm matrices in staphylococci. *Front Cell Infect Microbiol* 4:171. <https://doi.org/10.3389/fcimb.2014.00171>.
 49. Fong JNC, Yildiz FH. 3 April 2015. Biofilm matrix proteins. *Microbiol Spectr* <https://doi.org/10.1128/microbiolspec.MB-0004-2014>.
 50. Hobbey L, Harkins C, MacPhee CE, Stanley-Wall NR. 2015. Giving structure to the biofilm matrix: an overview of individual strategies and emerging common themes. *FEMS Microbiol Rev* 39:649–669. <https://doi.org/10.1093/femsre/fuv015>.
 51. Barbu EM, Mackenzie C, Foster TJ, Höök M. 2014. SdrC induces staphylococcal biofilm formation through a homophilic interaction. *Mol Microbiol* 94:172–185. <https://doi.org/10.1111/mmi.12750>.
 52. Feuillie C, Formosa-Dague C, Hays LMC, Vervaeck O, Derclaye S, Brennan MP, Foster TJ, Geoghegan JA, Dufrene YF. 2017. Molecular interactions and inhibition of the staphylococcal biofilm-forming protein SdrC. *Proc Natl Acad Sci U S A* 114:3738–3743. <https://doi.org/10.1073/pnas.1616805114>.
 53. Kalamajski S, Aspberg A, Lindblom K, Heinegård D, Oldberg A. 2009. Asporin competes with decorin for collagen binding, binds calcium and promotes osteoblast collagen mineralization. *Biochem J* 423:53–59. <https://doi.org/10.1042/BJ20090542>.
 54. Klänning E, Christensen B, Sørensen ES, Vorup-Jensen T, Jensen JK. 2014. Osteopontin binds multiple calcium ions with high affinity and independently of phosphorylation status. *Bone* 66:90–95. <https://doi.org/10.1016/j.bone.2014.05.020>.
 55. Isowa Y, Sarashina I, Setiamarga DHE, Endo K. 2012. A comparative study of the shell matrix protein aspein in pteridoid bivalves. *J Mol Evol* 75:11–18. <https://doi.org/10.1007/s00239-012-9514-3>.
 56. Decho AW. 2010. Overview of biopolymer-induced mineralization: what goes on in biofilms? *Ecol Eng* 36:137–144. <https://doi.org/10.1016/j.ecoleng.2009.01.003>.
 57. Takeuchi T, Sarashina I, Iijima M, Endo K. 2008. *In vitro* regulation of CaCO₃ crystal polymorphism by the highly acidic molluscan shell protein aspein. *FEBS Lett* 582:591–596. <https://doi.org/10.1016/j.febslet.2008.01.026>.
 58. Zhu T, Dittrich M. 2016. Carbonate precipitation through microbial activities in natural environment, and their potential in biotechnology: a review. *Front Bioeng Biotechnol* 4:4. <https://doi.org/10.3389/fbioe.2016.00004>.
 59. Bai Y, Guo XJ, Li YZ, Huang T. 2017. Experimental and visual research on the microbial induced carbonate precipitation by *Pseudomonas aeruginosa*. *AMB Express* 7:57. <https://doi.org/10.1186/s13568-017-0358-5>.
 60. Martínez-Gil M, Romero D, Kolter R, Espinosa-Urgel M. 2012. Calcium causes multimerization of the large adhesin LapF and modulates biofilm formation by *Pseudomonas putida*. *J Bacteriol* 194:6782–6789. <https://doi.org/10.1128/JB.01094-12>.
 61. de Vries GE, Harms N, Hoogendijk J, Stouthamer AH. 1989. Isolation and characterization of *Paracoccus denitrificans* mutants with increased conjugation frequencies and pleiotropic loss of a (nGATc) DNA-modifying property. *Arch Microbiol* 152:52–57. <https://doi.org/10.1007/BF00447011>.
 62. Simon R, Priefer U, Pühler A. 1983. A broad host range mobilization system for in vivo genetic engineering: transposon mutagenesis in Gram negative bacteria. *Biotechnology* 1:784–791. <https://doi.org/10.1038/nbt1183-784>.
 63. Shearer N, Hinsley AP, Van Spanning RJ, Spiro S. 1999. Anaerobic growth of *Paracoccus denitrificans* requires cobalamin: characterization of *cobK* and *cobJ* genes. *J Bacteriol* 181:6907–6913.
 64. Schäfer A, Tauch A, Jäger W, Kalinowski J, Thierbach G, Pühler A. 1994. Small mobilizable multi-purpose cloning vectors derived from the *Escherichia coli* plasmids pK18 and pK19: selection of defined deletions in the chromosome of *Corynebacterium glutamicum*. *Gene* 145:69–73. [https://doi.org/10.1016/0378-1119\(94\)90324-7](https://doi.org/10.1016/0378-1119(94)90324-7).
 65. Sullivan MJ, Gates AJ, Appia-Ayme C, Rowley G, Richardson DJ. 2013. Copper control of bacterial nitrous oxide emission and its impact on vitamin B12-dependent metabolism. *Proc Natl Acad Sci U S A* 110:19926–19931. <https://doi.org/10.1073/pnas.1314529110>.
 66. Wurch T, Lestienne F, Pauwels PJ. 1998. A modified overlap extension PCR method to create chimeric genes in the absence of restriction enzymes. *Biotechnol Tech* 12:653–657. <https://doi.org/10.1023/A:1008848517221>.
 67. Spaink HP, Okker RJH, Wijffelman CA, Pees E, Lugtenberg BJJ. 1987. Promoters in the nodulation region of the *Rhizobium leguminosarum* Sym plasmid pRL1J1. *Plant Mol Biol* 9:27–39. <https://doi.org/10.1007/BF00017984>.
 68. Pratt LA, Kolter R. 1998. Genetic analysis of *Escherichia coli* biofilm formation: roles of flagella, motility, chemotaxis and type I pili. *Mol Microbiol* 30:285–293. <https://doi.org/10.1046/j.1365-2958.1998.01061.x>.
 69. Miller JH. 1992. A short course in bacterial genetics. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
 70. Mehta HH, Liu Y, Zhang MQ, Spiro S. 2015. Genome-wide analysis of the response to nitric oxide in uropathogenic *Escherichia coli* CFT073. *Microb Genom* 1:e000031. <https://doi.org/10.1099/mgen.0.000031>.