

Supplementary Material

Coherent changes of cortical thickness and corresponding white matter microstructure during brain development accessed by diffusion MRI tractography from parcellated cortex

Tina Jeon^{1,2}, Virendra Mishra^{2,3}, Minhui Ouyang^{1,2}, Min Chen⁴, Hao Huang^{1,5}

¹Radiology Research, Children's Hospital of Philadelphia, Philadelphia, PA, USA

²Advanced Imaging Research Center, University of Texas Southwestern Medical Center at Dallas, Dallas, TX USA

³Lou Ruvo Center for Brain Health, Cleveland Clinic, Las Vegas, NV, USA

⁴Department of Mathematical Sciences, University of Texas at Dallas, Richardson, TX USA

⁵Department of Radiology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

***Correspondence:** Hao Huang, Ph.D., Radiology Research, Children's Hospital of Philadelphia, 3401 Civic Center Blvd, Philadelphia, PA, 19104, USA.

huangh6@email.chop.edu

Tina Jeon, Radiology Research, Children's Hospital of Philadelphia, 3401 Civic Center Blvd, Philadelphia, PA, 19104, USA.

jeont@email.chop.edu

Supplementary Figure 1: The histogram of the change slopes of cortical thickness (top panel) and corresponding white matter FA (bottom panel) for each of the 16 frontal lobe gyral regions during development from 7 to 25 years of age. The regions were grouped into 3 clusters, slow (green), median (yellow), and fast (red) based on slope values.

